

Methodology article

Open Access

A high-density collection of EMS-induced mutations for TILLING in Landsberg *erecta* genetic background of *Arabidopsis*

Beatriz Martín, Mercedes Ramiro, José M Martínez-Zapater and Carlos Alonso-Blanco*

Address: Departamento de Genética Molecular de Plantas, Centro Nacional de Biotecnología (CNB), Consejo Superior de Investigaciones Científicas (CSIC), Madrid-28049, Spain

Email: Beatriz Martín - bmartin@cnb.csic.es; Mercedes Ramiro - mramiro@cnb.csic.es; José M Martínez-Zapater - zapater@cnb.csic.es; Carlos Alonso-Blanco* - calonso@cnb.csic.es

* Corresponding author

Published: 14 December 2009

Received: 6 July 2009

BMC Plant Biology 2009, **9**:147 doi:10.1186/1471-2229-9-147

Accepted: 14 December 2009

This article is available from: <http://www.biomedcentral.com/1471-2229/9/147>

© 2009 Martín et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: *Arabidopsis thaliana* is the main model species for plant molecular genetics studies and world-wide efforts are devoted to identify the function of all its genes. To this end, reverse genetics by TILLING (Targeting Induced Local Lesions IN Genomes) in a permanent collection of chemically induced mutants is providing a unique resource in Columbia genetic background. In this work, we aim to extend TILLING resources available in *A. thaliana* by developing a new population of ethyl methanesulphonate (EMS) induced mutants in the second commonest reference strain. In addition, we pursue to saturate the number of EMS induced mutations that can be tolerated by viable and fertile plants.

Results: By mutagenizing with different EMS concentrations we have developed a permanent collection of 3712 M₂/M₃ independent mutant lines in the reference strain Landsberg *erecta* (Ler) of *A. thaliana*. This population has been named as the Arabidopsis TILLer collection. The frequency of mutations per line was maximized by using M₁ plants with low but sufficient seed fertility. Application of TILLING to search for mutants in 14 genes identified 21 to 46 mutations per gene, which correspond to a total of 450 mutations. Missense mutations were found for all genes while truncations were selected for all except one. We estimated that, on average, these lines carry one mutation every 89 kb, Ler population providing a total of more than five million induced mutations. It is estimated that TILLer collection shows a two to three fold higher EMS mutation density per individual than previously reported *A. thaliana* population.

Conclusions: Analysis of TILLer collection demonstrates its usefulness for large scale TILLING reverse genetics in another reference genetic background of *A. thaliana*. Comparisons with TILLING populations in other organisms indicate that this new *A. thaliana* collection carries the highest chemically induced mutation density per individual known in diploid species.

Background

A major challenge in plant biology is the identification of biological functions for all genes from the main model plant species, *Arabidopsis thaliana* and rice. To this end, a large number of genetics and genomics resources are being developed in both model plants [1,2]. In particular, collections of induced mutants that can be screened by reverse genetics, such as T-DNA or transposon insertional mutants [3-5] provide a unique resource for functional studies. However, the mutational spectrum of insertional mutagenesis with effect on gene function is mostly limited to gene knock-out disruptions. Genes whose severe loss-of-function is lethal or highly pleiotropic cannot be functionally dissected with such mutants. In addition, the size of saturated populations containing insertion mutants randomly generated for most genes of an organism is extremely high because each line carries only a rather small number of mutations [6]. As a complementary resource, chemically induced mutants have been shown to provide an efficient alternative because each individual line can bear single point missense and nonsense substitutions in hundreds of genes [7]. Therefore, an allelic series of induced mutations with different effects on gene function can be easily isolated by screening a few thousands mutagenized plants [6].

In the past few years, chemically induced mutants have become a major resource for reverse genetics studies thanks to the development of TILLING (Targeting Induced Local Lesions IN Genomes) [8]. TILLING enables the reverse selection of single point mutations by cleavage of mismatches in heteroduplex DNA with the endonuclease CEL I. This powerful strategy was first applied in an *A. thaliana* mutant collection induced with ethyl methanesulphonate (EMS) [9,10] in the commonest genetic background Columbia (Col) whose genome sequence had been first completed [11]. Since then, TILLING collections of EMS induced mutants have been developed in a large number of plant species including rice, maize, barley, sorghum, wheat, *Brassica napus*, *B. oleracea* and *Medicago truncatula*, as well as model animals like *Drosophila* and *Caenorhabditis elegans* [12-21]. In most of these EMS mutant collections, reference genetic backgrounds of wide and general interest are used. However, given the limitations of having mutations in a single genetic background, new populations of chemically induced mutants for TILLING analyses are currently being developed in other reference strains of several species like rice or soybean [22,23]. In addition, the quality of TILLING mutant populations is determined by the density of mutations per individual, since this limits the size of allelic series than can be isolated for each gene and the size of a saturated genome population. For this reason, other TILLING populations have been developed in rice, barley, soybean or *M. truncatula*, aiming to increase the amount of mutations per line

by either using different mutagens like sodium azide and N-methyl-N-nitrosourea or increasing the mutagen dose [12,22-25].

In *A. thaliana*, several reference genetic backgrounds are widely used such as Col or Landsberg *erecta* (*Ler*). The latter is the second most commonly studied strain because many mutants have been classically isolated in it and a large portion of its genome sequence was available soon after Col sequence [26]. In this work we have developed a new collection of *A. thaliana* EMS induced mutants for TILLING reverse genetics, aiming at two major objectives. First, to extend TILLING resources in *A. thaliana* by using *Ler* reference genetic background, for which reverse genetic tools are rather limited. Second, to enrich the number of independent mutations available in this collection as much as possible by increasing the density of mutations per line. TILLING evaluation of this population for several gene fragments indicates that it carries the largest density of chemically induced mutations reported in diploid organisms, hence demonstrating its usefulness for reverse selection of mutants.

Results

Generation of a permanent collection of highly EMS-mutagenized lines in *Arabidopsis*

To obtain a new population of chemically induced mutant lines useful for reverse genetic studies in *Arabidopsis thaliana*, seeds of the Landsberg *erecta* (*Ler*) *glabrous1-1* genotype were mutagenized with EMS at concentrations of 20 to 50 mM (Figure 1). The effects of EMS and the efficiency of the mutagenesis treatment were estimated by quantifying three parameters on M_1 plants: seed germination, frequency of albino chimeras and fertility (see Methods). Germination of M_1 seeds was negatively correlated with EMS dose ($r = -0.93$; $p = 0.008$), while the frequency of M_1 albino chimeras increased with concentration ($r = 0.99$; $p = 0.001$) (Figure 1A and 1B). Seed fertility of M_1 plants and the degree of M_2 embryo lethality was quantified by estimating the proportion of fully or nearly sterile fruits (classes As and Aa) and the proportion of semi or normal fertile fruits (classes B and C). As shown in Figure 1C, the total frequency of class A fruits increased linearly with EMS concentration, whereas the frequency of fertile fruits rapidly decreased. To maximize the frequency of mutations per individual, only M_1 plants from treatments showing a frequency of fertile fruits smaller than 35% but larger than 2% were individually harvested. A total of 3712 M_2 families were grown to isolate individual M_2 DNA and to harvest their M_3 offspring seeds. To ensure independence of the mutations present in this population, a single M_2 plant was harvested from each M_1 plant. In agreement with the high proportion of embryo lethality, all M_2 families segregated for easily visible morphological mutations (data not shown). Fifty six percent

Table 1: Description of *A. thaliana* Ler mutant lines and mutations in relation to EMS dose.

EMS dose (mM)	Number of TILLer lines	Mean (B+C) fertility class (%)	Total number of screened lines*	Number of mutations	Het/Hom mutation ratio	Number of lines with two mutations	Density of mutations per line
20	46	24.5 ± 5.4	614 (1.5%)	7 (1.5%)	6.0	0 (0%)	1/84 kb
25	2082	16.8 ± 3.7	21150 (50.8%)	178 (39.6%)	3.6	10 (29.4%)	1/114 kb
30	741	10.5 ± 2.8	9196 (22.1%)	112 (24.9%)	3.3	10 (29.4%)	1/79 kb
35	739	3.7 ± 1.2	9313 (22.4%)	128 (28.4%)	4.3	12 (35.3%)	1/70 kb
40	104	2.6 ± 1.0	1335 (3.2%)	25 (5.6%)	4.0	2 (5.9%)	1/51 kb
Total	3712	12.6 ± 4.6**	41608	450	3.7**	34	1/96 kb**

*: Number of screened lines is estimated as the total sum of individuals analyzed for each of the 14 gene fragments.

** : Weighted averages are shown according to frequency of lines of the different EMS dose.

of M_2 lines were derived from 25 mM EMS mutagenesis, and on average, EMS treatments used to obtain the collection show less than 25% fertile fruits (Table 1). The DNA of M_2 plants and the M_3 seeds of the 3712 lines were stored (see Methods) providing a permanent population of mutant lines for TILLING analysis in *Ler* genetic background. This population has been named as the Arabidopsis TILLer collection.

Mutation frequency, distribution and functional spectrum in TILLer collection

The quality of this collection for mutant discovery was evaluated after the analysis of 14 gene fragments distributed among four of the five *A. thaliana* chromosomes and chosen from requests by different research laboratories (see Methods; Figure 2 and Table 2). These fragments have a GC content similar to that of fragments studied in Col background TILLING collection [10] and to average coding regions of *A. thaliana* genome [11]. On average, amplicons were 1.1 kb long and 62.2% corresponded to exon sequence, which is similar to genome average exon proportion [11]. In total, we found and confirmed by sequencing 450 mutations in the 15.7 kb analyzed from different amplicons. All mutations corresponded to G/C to A/T transitions, in agreement with the nearly unique type of nucleotide substitutions observed in previous analyses of EMS-mutagenized *A. thaliana* plants [10]. The distribution of mutations among the EMS doses of the lines was independent of the number of lines per dose (Table 1; $\chi^2 = 28.9$; $df = 4$; $p < 0.0001$). A larger number of mutations were found at 30, 35 and 40 mM than expected from the number of lines, while the opposite was found at 25 mM. In addition, 34 lines carried several mutations in the same or two different gene fragments (Table 1). An excess of these lines was found in plants derived from high EMS concentrations (≥ 30 mM) while a deficiency appears in low EMS concentrations (≤ 25 mM) when comparing with the expected number according to the proportion of lines from each EMS dose ($\chi^2 = 5.8$; $df = 1$; $p = 0.016$). Therefore, the higher the EMS concentration used

to obtain the lines, the larger the number of mutations per line.

On average we analyzed 2972 lines per fragment and detected 10.8 mutations per 1000 mutant lines (Table 2). Twenty-one to 46 mutations were found per fragment, and in most gene fragments there was a reduction of mutation detection in the ~ 100 bp terminal segments (Figure 2), as expected from LI-COR detection system (see Methods). However, mutations appeared evenly distributed along the rest of the gene fragments within exon and intron regions (Figure 2).

For all but one gene fragment we found mutations of three classes according to their predicted effects on protein structure: silent, missense and truncation mutations (Table 2). The observed frequencies of the three classes of mutations fitted the expected frequencies of silent, missense and truncations, respectively, as estimated by CODDLE analyses ($\chi^2 = 1.7$; $df = 2$; $p = 0.42$). Truncations include nonsense mutations generating premature stop codons and mutations in intron splice sites, the observed frequencies of both classes (2.5% and 1.8% respectively) also fitting expected frequencies (4.0% and 1.1%) ($\chi^2 = 2.8$; $df = 1$; $p = 0.09$). Interestingly, truncation mutations were obtained for 13 of the 14 fragments, as expected from their 5.1% frequency and the large average number of mutations found per gene ($1 - [1 - 0.05]^{32} = 0.81$ probability).

As shown in Table 2, an average ratio of heterozygous/homozygous mutations of 3.7 was found, which is significantly different from the expected 2:1 proportion for M_2 plants ($\chi^2 = 30.2$; $df = 1$; $p < 0.0001$). Although an excess of heterozygotes appeared for silent mutations ($p < 0.01$), this ratio was extreme for truncations since all but one of such mutations were present as heterozygotes. In addition, distortion from the expected proportion was larger for high EMS dose lines (35-40 mM) than for low concentrations (25-30 mM) (Table 1).




Figure 1
Dose effects of EMS mutagenesis in *A. thaliana* M₁ Ler plants. A) Frequency of M₁ seed germination. B) Frequency of M₁ albino chimeras at vegetative stage. C) Frequency of sterility, embryo lethality and fertility measured as the percentage of different fruit classes. Fruits of M₁ plants are classified as As, Aa, B and C (from fully sterile to nearly normal fertility) depending on the proportion of aborted seeds (see Methods for details). Data are mean ± SE of three to six replicates. Untreated control plants showed 100% germination, 0% albino chimeras and 0, 3 and 97% of A, B and C fruit classes.

From these analyses we estimated an average density of detected mutations per line of 1 mutation per 89 kb (450 mutations/[2972 lines × 13.4 kb]), which was calculated after subtracting 160 terminal base pairs with low LI-COR detection, from each amplicon (see Methods). However, the density of mutations varied from 1/114 kb to 1/51 kb depending on the EMS dose used to generate the lines, a two-fold variation being found between 25 and 40 mM (Table 1). To contrast this average mutation frequency estimation, the density of mutations was also independently calculated from the number of pool samples with two mutant individuals in the same fragment or from the number of individual lines with two mutations in the same gene fragment [10]. Forty-four pool samples were found to carry two mutant individuals when analyzing the individual lines. Thus, a total of 406 pool samples were originally detected as positive pools, which contain 406 × 8 individuals representing a sample analyzed at individual level to find second mutations. From these 406 pool samples with at least one positive line we estimated a density of 1 mutation/71 kb (44/[406 pool samples × 8 individuals × 0.96 kb]), which is similar to previous estimate. On the other hand, when sequencing positive lines for their corresponding fragments, five individual lines were found to carry two mutations within the same fragment. Therefore, 445 lines were sequenced and can be considered a sample analyzed to detect second mutations by sequencing. From these lines we calculated a density of 1 mutation/100 kb (5/[445 × 1.12 kb]), which is comparable to the above estimates. In contrast to previous calculations, this latest density was estimated from the complete amplicon length (1.12 kb) because it was derived from the sequencing of entire fragments and not from LI-COR detection of positive lines.

From the above density of mutations we have calculated an average number of 1404 mutations per line, the complete TILLer collection providing a total of 5.2 million mutations. Taking into account the observed frequencies of truncation and missense mutations (Table 2), and the total length of gene regions of *A. thaliana* genome (see Methods), we have roughly estimated that each TILLer line contains, on average, 30 genes with knock-out mutations and 281 genes with aminoacid substitutions.

Discussion

We have developed a new permanent collection of 3712 independent EMS-induced mutant lines for reverse genetic analysis in the reference laboratory strain Landsberg *erecta* of *A. thaliana*. To maximize the number of mutations present in this population we have increased the frequency of mutations per M₂/M₃ line by using M₁ plants with lower seed fertility than that of plants used to obtain the existing population in Columbia background [9]. By compromising fertility, we aimed to saturate the

Table 2: Mutations found in 14 gene fragments analyzed in TILLer collection.

Gene	Amplicon length (bp)	Exon length (bp)	Intron length (bp)	GC content (%)	# of screened lines	Total # of mutations	Mutation class*		
							Silent	Missense	Truncation**
At1g12650	977	416	561	35.6	1968	22	11/1	6/3	1/0
At1g32640	1015	1015	0	49.8	2000	42	8/2	22/7	3/0
At2g22475	839	439	400	44.3	3416	33	7/2	16/7	1/0
At2g22540	1128	278	850	35.5	3216	25	18/2	4/0	1/0
At2g26300	1152	547	605	37.7	3456	32	19/3	6/2	2/0
At2g27100	1272	1049	223	43.8	3360	25	6/2	14/3	0/0
At3g06120	1143	513	630	35.0	3432	29	11/3	10/4	1/0
At3g13040	1093	581	512	39.8	2728	28	17/3	4/2	2/0
At3g24740	1195	1068	127	46.4	3472	46	10/3	23/8	2/0
At5g07280	1228	1228	0	46.0	2728	36	6/2	17/10	1/0
At5g11270	1075	667	408	39.8	1896	21	7/5	4/4	1/0
At5g23280	1077	762	315	42.5	3352	39	7/1	23/7	1/0
At5g41560	1299	262	1037	36.8	3200	32	25/5	0/1	1/0
At5g50570	1180	939	241	40.3	3384	40	12/1	23/1	2/1
Total	15673	9764	5909	41.0	41608	450	164/35	172/59	19/1
Average	1120	697	422	41.0	2972	32	12/3	12/4	1/0
Observed frequency (%)							47.2	48.5	4.3
Expected frequency (%)							48.8	46.1	5.1

*: number of heterozygous/homozygous mutations are given for each of the three mutations classes.

***: Truncations include mutations generating premature stop codons and mutations in splice sites.

number of chemically induced mutations that can be tolerated by *A. thaliana* plants that are still viable and able of sexual reproduction. We estimated that, on average, the lines of this new *Ler* collection carry one mutation every 89 kb, which is significantly larger than the density of 1/300 kb estimated in current Col population [10]. As expected, we found that the higher the EMS concentration the higher the density of detected mutations per line. Thus, experimental control of EMS mutagenesis enables substantial increase of the frequency of induced mutations in viable and seed fertile plants. However, we cannot discard that mutation density differences between both TILLING populations of *A. thaliana* might be partly due to natural genetic variation between both wild type strains for their tolerance to chemically induced mutations. Accordingly, it could be speculated that such natural variation might be determined by variation for reproductive system plasticity or for DNA repair mechanisms.

As described by Greene *et al.* [10] estimations of the density of chemically induced mutations detected by TILLING procedure can be biased due to several factors such as: 1) uneven mutation detection among the pools of eight plants; 2) uneven mutation detection along the length of gene fragments; 3) higher GC content of analyzed frag-

ments (41%) than average genome (35%) [11]; and 4) dilution of one fourth of the M_1 mutations in M_2 plants. Taking into account these factors, it has been estimated that Col population shows a corrected average mutation density per line of 1/170 kb. Despite these factors, the 1/89 kb average frequency of induced mutations detected in *Ler* population is nearly twice higher than that of Col. Furthermore, *Ler* lines obtained from the highest EMS dose show a 1/51 kb mutation density, which triplicates the lowest density estimate of Col population. Comparisons with TILLING populations in other species indicates that *A. thaliana Ler* collection carries the highest mutation density induced by different chemical agents in diploid plants, including *Brassica oleracea*, *Medicago truncatula*, rice, barley, maize and sorghum; or animals like *Drosophila* and *Caenorhabditis elegans*, or the oomycote *Phytophthora sojae* (see Table 3). Only rapeseed and wheat collections carry a higher density of mutations, as expected from their polyploid nature and consequently, their higher tolerance to loss-of-function mutations due to gene duplications and redundancies (Table 3). Thus, the density of mutations found in *Ler A. thaliana* population increases the estimated load of chemically induced mutations that diploid species can tolerate in sexually fertile individuals.




Figure 2
Gene fragments analyzed and mutations found in TILLer collection. Positions of silent, missense, splicing site and nonsense mutations are indicated by grey arrow heads, black arrow heads, grey asterisks and black asterisks, respectively.

The two *A. thaliana* TILLING populations, *Ler* and *Col*, also differ in the proportion of heterozygous:homozygous mutations recovered in TILLING analyses, *Ler* showing substantially higher total average ratio than *Col* (3.7 versus 2.1, respectively) [10]. The largest deficiency of homozygous mutations corresponds to truncations, which shows the largest difference between both populations (ratio of 3.7 vs. 19 for *Col* and *Ler*, respectively). Therefore, a stronger negative selection against deleterious mutations seems to affect *Ler* than *Col* collection. This is probably a consequence of the extreme high-density of mutations present in *Ler* lines, since the maximum load of deleterious induced mutations that can be tolerated by a viable and fertile M_2 plant will likely be determined by a threshold number of homozygous truncations and deleterious missense mutations. M_2 plants carrying a higher number of homozygous deleterious mutations than this threshold will not be viable or fertile. Given the self-fertilizing nature of *A. thaliana*, the higher the M_1 mutation density, the higher the proportion of M_2 offspring plants that will surpass the maximum number of homozygous deleterious mutations and, consequently, stronger selection against such mutations. Thus, higher M_1 mutation densities will lead to higher M_2 ratios of heterozygous/homozygous mutations due to lower frequency of M_2 plants below the threshold of homozygous deleterious mutations. This relationship is supported by the larger ratios observed in *Ler* lines with high mutation density generated with EMS doses ≥ 35 mM, than in lines with lower density obtained with 25-30 mM EMS. Nevertheless, presumed silent mutations including synonymous and intronic mutations also showed a significant defect of homozygotes in *Ler* collection, whereas this was not observed for missense mutations. Potential genetic mechanisms accounting for this unexpected result are unknown but it cannot be discarded that the genes surveyed in this work are biased for the deleterious effect of their mutations. In agreement, other *A. thaliana* public mutant collections do not contain mutations in several of the genes analyzed here <http://www.arabidopsis.org> suggesting that mutations in their coding and non-coding regions show stronger deleterious defects than genome average.

Conclusions

The TILLer collection generated in this work provides a new resource for reverse selection of EMS induced mutants in *A. thaliana*. The high mutation density of this population increases the size of allelic series that can be obtained and reduces the population size that needs to be screened. However, this high mutation density implies that more backcrosses are required to eliminate undesired background mutations in selected mutant lines. It has been estimated that 20 mutations are necessary to have 0.95 probability of finding a missense deleterious mutation per gene [6]. Considering the ~50% observed fre-

Table 3: Frequency of chemically induced mutations reported by TILLING in different species.

Species	Ploidy level	Mutagen	Mutation density per line	Reference
<i>A. thaliana</i> (Landsberg erecta)	2x	EMS	1/89 kb	present work
<i>A. thaliana</i> (Columbia)	2x	EMS	1/300 kb	[10]
<i>Brassica oleracea</i>	2x	EMS	1/447 kb	[18]
<i>Brassica napus</i>	4x	EMS	1/41.5 kb	[17]
<i>Medicago truncatula</i>	2x	EMS	1/400 kb	[19]
<i>Medicago truncatula</i>	2x	EMS	1/485 kb	[25]
<i>Glycine max</i>	4x	EMS	1/140 to 1/550 kb	[23]
<i>Glycine max</i>	4x	NMU	1/140 kb	[23]
<i>Oryza sativa</i>	2x	EMS	1/294 kb	[12]
<i>Oryza sativa</i>	2x	NaN ₃ + MNU	1/265 kb	[12]
<i>Oryza sativa</i>	2x	MNU	1/135 kb	[22]
<i>Zea mays</i>	2x	EMS	1/485 kb	[13]
<i>Sorghum bicolor</i>	2x	EMS	1/526 kb	[15]
<i>Hordeum vulgare</i>	2x	EMS	1/1000 kb	[14]
<i>Hordeum vulgare</i>	2x	NaN ₃	1/374 kb	[24]
<i>Triticum aestivum</i>	6x	EMS	1/40 kb	[16]
<i>Triticum durum</i>	4x	EMS	1/25 kb	[16]
<i>Caenorhabditis elegans</i>	2x	EMS	1/293 kb	[21]
<i>Drosophila melanogaster</i>	2x	EMS	1/91 to 1/156 kb	[20]
<i>Phytophthora sojae</i>	2x	ENU	1/142 kb	[30]

NaN₃: sodium azide; MNU: N-methyl-N-nitrosourea; ENU: ethylnitrosourea.

quency of missense mutations and assuming that 25% of them are deleterious, we have calculated that on average, 1774 TILLer lines are sufficient to obtain 20 mutations per ~1 kb gene fragment, while the larger analyses carried out until now are providing additional truncation mutations for ~90% of the genes. Currently, TILLer collection is screened as a public service to search for mutants in genes of interest for any laboratory <http://www.cnb.csic.es/~tiller/>. The availability of another TILLING service in the second commonest reference genetic background of *A. thaliana* enables deeper gene functional studies such as those aiming to uncover new gene effects or interactions of specific mutations with genetic backgrounds. Given the success of current existing collections, it can be expected that the use of chemically induced genetic variation will further extend in the near future with the development of similar resources in other reference strains and/or using other mutagens.

Methods

Mutagenesis

Seeds of the laboratory strain Landsberg *erecta* carrying the marker mutation *glabrous1-1* were mutagenized with ethyl methanesulphonate (EMS) [9]. Fresh M₀ seeds were treated with 20, 25, 30, 35, 40 or 50 mM EMS during 17 hours in 10 ml vials containing 2500 seeds. Three to eight batches of 2500 seeds (vials) were treated at each dose. After thorough washing, M₁ seeds were sown on pots with soil:vermiculite mix at 3:1 proportion, in a 20°C greenhouse supplemented with lamps to provide a 16 hours light:8 hours darkness photoperiod. To estimate the EMS effects and the quality of the mutagenesis we quantified

germination of M₁ seeds, and the proportion of chimeric M₁ plants that show albino or yellow sectors at the vegetative stage of six-eight leaves (albino chimeras). In addition, seed fertility and degree of embryo lethality of the M₁ plants were estimated as previously described [9] with the following modifications. For each EMS dose, 10 mature siliques of the main inflorescence from 10 M₁ plants were dissected under a stereomicroscope and the number of normal and aborted seeds was counted. From these data, fruits were classified in four classes according to their proportion of normal M₂ seeds and M₂ defective embryos. Class As is completely sterile and has no seed, either normal or aborted; class Aa has a 3:1 proportion of normal:aborted seeds, or smaller (aborted seeds >20%); class B shows 4:1 to 13:1 proportions (20% ≥ aborted seeds >6.7%); and class C has 14:1 or larger proportion (nearly normal fertile fruits with less than 6.7% aborted seeds). M₁ plants were individually harvested and treatments with a frequency of fertile fruits (B+C) larger than 2% or smaller than 35% were used to generate the M₂/M₃ lines that are part of TILLer collection (Table 1). Mutagenized batches with more than 35% or less than 2% fertile fruits were discarded independently of the concentration of their EMS dose.

Development of mutant and DNA collections

Five to sixteen M₂ offspring seedlings were grown from each M₁ individual and tissue was collected from a single M₂ fertile plant. M₃ seeds of each M₂ selected plant were individually harvested and stored. To ensure enough tissue and M₃ offspring seeds from a single M₂ plant, each family was grown on a 0.9 l. pot that was maintained until

fruit formation in a growth chamber illuminated with cool-white fluorescent lamps that provide a short day photoperiod of 8 hours light:16 hours darkness. DNA was isolated as previously described [27] without mercaptoethanol. The DNA of 3712 M₂ plants was quantified, diluted and arrayed in a total of 58 (8 × 8)-individual plates as described in [28]. The DNA was combined in groups of eight individuals using a one-dimension pooling strategy. Thus, the 3712 samples of the collection were arranged in five (96 × 8)-pool plates, four containing the DNA of 768 individuals and one from 640 individuals.

Mutation detection

Mutations in gene fragments were detected using the TILLING procedure developed and described by Till *et al.* [9,28]. Briefly, primers for amplification of target genes were designed using the CODDLE and Primer 3 system <http://blocks.fhcrc.org/proweb/input/>. Forward and reverse primers were labeled with IRDye 700 and IRDye 800 respectively. PCR, heteroduplex DNA formation and heteroduplex digestion with CEL I were carried out using the 768 pool plates containing 5 µl of DNA at 0.15 ng/µl as previously described [28]. CEL I was purified from celery juice extracts according to [29] with minor modifications. For that, concentrated extracts were incubated with concanavalin A-sepharose, followed by chromatography purification steps with DEAE FF and Q columns using an AKTA FPLC system (Amersham). Cleaved DNA fragments were separated in a LI-COR 4300 DNA analyzer and gel images were manually analyzed using Photoshop (Adobe system) to find positive pools. Thereafter, individual plates containing the eight individual DNA samples of each positive pool were similarly analyzed after mixing with control wild type DNA. Validated mutants were sequenced and sequences were analyzed with Chromas and DNASTAR softwares. Mutation frequencies were calculated as described in [10] by subtracting 160 bp from the size of each amplicon due to the observed low ability of LI-COR system for mutation detection in the 80 bp terminal segments of gene fragments. To enable direct comparisons of mutation frequencies among TILLING projects from different species [10,12-25,30], these calculations were based on haploid sizes of the analyzed fragments. Thus, frequencies are given per kb of diploid genome and should be divided by two when taking into account the diploid nature of *A. thaliana*. The total number of mutations was calculated using an *A. thaliana* genome size of 125 Mb and a total size of gene regions of 33249 kb (exons) plus 18055 kb (introns) [11].

Authors' contributions

BM participated in the generation of the mutant collection, in the analysis of mutations and in the writing of the manuscript. MR participated in the generation of the mutant collection and in the analysis of mutations. JMZ

participated in the conception and design of the study and in the writing of manuscript. CAB participated in the conception, design and supervision of the study, in the analysis of mutations and in the writing of manuscript. All authors read and approved the final manuscript.

Acknowledgements

We thank Brad Till and the Arabidopsis TILLING project for kind assistance in setting up the TILLING procedure, and Leonor Kremer, Otto Törjék and Thomas Altmann for help with CEL I purification. This work and service has been funded by grants GEN200-4890-C07-01 and GEN2006-28555-EI from the Ministerio de Ciencia e Innovación of Spain.

References

- Borevitz JO, Ecker JR: **Plant genomics: the third wave.** *Annu Rev Genomics Hum Genet* 2004, **5**:443-477.
- Sasaki T, Matsumoto T, Antonio BA, Nagamura Y: **From mapping to sequencing, post-sequencing and beyond.** *Plant Cell Physiol* 2005, **46**(1):3-13.
- Alonso JM, Stepanova AN, Leisse TJ, Kim CJ, Chen H, Shinn P, Stevenson DK, Zimmerman J, Barajas P, Cheuk R, *et al.*: **Genome-wide insertional mutagenesis of *Arabidopsis thaliana*.** *Science* 2003, **301**(5633):653-657.
- Rosso MG, Li Y, Strizhov N, Reiss B, Dekker K, Weisshaar B: **An *Arabidopsis thaliana* T-DNA mutagenized population (GABI-Kat) for flanking sequence tag-based reverse genetics.** *Plant Mol Biol* 2003, **53**(1-2):247-259.
- An G, Jeong DH, Jung KH, Lee S: **Reverse genetic approaches for functional genomics of rice.** *Plant Mol Biol* 2005, **59**(1):111-123.
- Østergaard L, Yanofsky MF: **Establishing gene function by mutagenesis in *Arabidopsis thaliana*.** *Plant J* 2004, **39**(5):682-696.
- Henikoff S, Till BJ, Comai L: **TILLING. Traditional mutagenesis meets functional genomics.** *Plant Physiol* 2004, **135**(2):630-636.
- Colbert T, Till BJ, Tompa R, Reynolds S, Steine MN, Yeung AT, McCallum CM, Comai L, Henikoff S: **High-throughput screening for induced point mutations.** *Plant Physiol* 2001, **126**(2):480-484.
- Till BJ, Reynolds SH, Greene EA, Codomo CA, Enns LC, Johnson JE, Burtner C, Odden AR, Young K, Taylor NE, *et al.*: **Large-scale discovery of induced point mutations with high-throughput TILLING.** *Genome Res* 2003, **13**(3):524-530.
- Greene EA, Codomo CA, Taylor NE, Henikoff JG, Till BJ, Reynolds SH, Enns LC, Burtner C, Johnson JE, Odden AR, *et al.*: **Spectrum of chemically induced mutations from a large-scale reverse-genetic screen in *Arabidopsis*.** *Genetics* 2003, **164**(2):731-740.
- AGI: **Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*.** *Nature* 2000, **408**(6814):796-815.
- Till BJ, Cooper J, Tai TH, Colowit P, Greene EA, Henikoff S, Comai L: **Discovery of chemically induced mutations in rice by TILLING.** *BMC Plant Biol* 2007, **7**:19.
- Till BJ, Reynolds SH, Weil C, Springer N, Burtner C, Young K, Bowers E, Codomo CA, Enns LC, Odden AR, *et al.*: **Discovery of induced mutations in maize genes by TILLING.** *BMC Plant Biol* 2004, **4**:12.
- Caldwell DG, McCallum N, Shaw P, Muehlbauer GJ, Marshall DF, Waugh R: **A structured mutant population for forward and reverse genetics in barley (*Hordeum vulgare* L.).** *Plant J* 2004, **40**(1):143-150.
- Xin Z, Wang ML, Barkley NA, Burrow G, Franks C, Pederson G, Burke J: **Applying genotyping (TILLING) and phenotyping analyses to elucidate gene function in a chemically induced sorghum mutant population.** *BMC Plant Biol* 2008, **8**:103.
- Slade AJ, Fuerstenberg SI, Loeffler D, Steine MN, Facciotti D: **A reverse genetic, nontransgenic approach to wheat crop improvement by TILLING.** *Nat Biotechnol* 2005, **23**(1):75-81.
- Wang N, Wang Y, Tian F, King GJ, Zhang C, Long Y, Shi L, Meng J: **A functional genomics resource for *Brassica napus*: development of an EMS mutagenized population and discovery of FAEI point mutations by TILLING.** *New Phytol* 2008, **180**(4):751-765.
- Himelblau E, Gilchrist EJ, Buono K, Bizzell C, Mentzer L, Vogelzang R, Osborn T, Amasino RM, Parkin IA, Haughn GW: **Forward and**

- reverse genetics of rapid-cycling *Brassica oleracea*.** *Theor Appl Genet* 2009, **118(5)**:953-961.
19. Porceddu A, Panara F, Calderini O, Molinari L, Taviani P, Lanfaloni L, Scotti C, Carelli M, Scaramelli L, Bruschi G, et al.: **An Italian functional genomic resource for *Medicago truncatula*.** *BMC Res Notes* 2008, **1**:129.
 20. Winkler S, Schwabedissen A, Backasch D, Bokel C, Seidel C, Bonisch S, Furthauer M, Kuhrs A, Cobreros L, Brand M, et al.: **Target-selected mutant screen by TILLING in *Drosophila*.** *Genome Res* 2005, **15(5)**:718-723.
 21. Gilchrist EJ, O'Neil NJ, Rose AM, Zetka MC, Haughn GW: **TILLING is an effective reverse genetics technique for *Caenorhabditis elegans*.** *BMC Genomics* 2006, **7**:262.
 22. Suzuki T, Eiguchi M, Kumamaru T, Satoh H, Matsusaka H, Moriguchi K, Nagato Y, Kurata N: **MNU-induced mutant pools and high performance TILLING enable finding of any gene mutation in rice.** *Mol Genet Genomics* 2008, **279(3)**:213-223.
 23. Cooper JL, Till BJ, Laport RG, Darlow MC, Kleffner JM, Jamai A, El-Mellouki T, Liu S, Ritchie R, Nielsen N, et al.: **TILLING to detect induced mutations in soybean.** *BMC Plant Biol* 2008, **8**:9.
 24. Talamè V, Bovina R, Sanguineti MC, Tuberosa R, Lundqvist U, Salvi S: **TILLMore, a resource for the discovery of chemically induced mutants in barley.** *Plant Biotechnol J* 2008, **6(5)**:477-485.
 25. Le Signor C, Savoie V, Aubert G, Verdier J, Nicolas M, Pagny G, Moussy F, Sanchez M, Baker D, Clarke J, et al.: **Optimizing TILLING populations for reverse genetics in *Medicago truncatula*.** *Plant Biotechnol J* 2009, **7(5)**:430-441.
 26. Jander G, Norris SR, Rounsley SD, Bush DF, Levin IM, Last RL: **Ara-bidopsis map-based cloning in the post-genome era.** *Plant Physiol* 2002, **129(2)**:440-450.
 27. Bernartzky R, Tanksley S: **Genetics of acting-related sequences in tomato.** *Theor Appl Genet* 1986, **72**:314-324.
 28. Till BJ, Zerr T, Comai L, Henikoff S: **A protocol for TILLING and Ecotilling in plants and animals.** *Nat Protoc* 2006, **1(5)**:2465-2477.
 29. Oleykowski CA, Bronson Mullins CR, Godwin AK, Yeung AT: **Mutation detection using a novel plant endonuclease.** *Nucleic Acids Res* 1998, **26(20)**:4597-4602.
 30. Lamour KH, Finley L, Hurtado-Gonzales O, Gobena D, Tierney M, Meijer HJ: **Targeted gene mutation in *Phytophthora spp.*** *Mol Plant Microbe Interact* 2006, **19(12)**:1359-1367.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

