

CORRECTION

Open Access



Correction: Genome-wide characterization and identification of candidate *ERF* genes involved in various abiotic stress responses in sesame (*Sesamum indicum* L.)

Ruqi Su^{1,2}, Senouwa Segla Koffi Dossou², Komivi Dossa^{3,4}, Rong Zhou², Aili Liu², Yanping Zhong^{1,2}, Sheng Fang¹, Xiurong Zhang², Ziming Wu^{1*} and Jun You^{2*}

Correction: BMC Plant Biol 22, 256 (2022)

<https://doi.org/10.1186/s12870-022-03632-7>

Following publication of the original article [1], authors identified an error found in Affiliations 1 and 2. The correct affiliation details are presented below:

¹Key Laboratory of Crop Physiology, Ecology, and Genetic Breeding, Ministry of Education/College of Agronomy, Jiangxi Agricultural University, Nanchang 330045, China

²Key Laboratory of Biology and Genetic Improvement of Oil Crops of the Ministry of Agriculture and Rural Affairs, Oil Crops Research Institute of the Chinese Academy of Agricultural Sciences, Wuhan 430062, China

The corrections do not have any effect on the final conclusions of the paper. The original article has been corrected.

Author details

¹Key Laboratory of Crop Physiology, Ecology, and Genetic Breeding, Ministry of Education/College of Agronomy, Jiangxi Agricultural University, Nanchang 330045, China. ²Key Laboratory of Biology and Genetic Improvement of Oil Crops of the Ministry of Agriculture and Rural Affairs, Oil Crops Research Institute of the Chinese Academy of Agricultural Sciences, Wuhan 430062, China. ³CIRAD, UMR AGAP Institut, F-34398 Montpellier, France. ⁴UMR AGAP Institut, Univ Montpellier, CIRAD, INRAE, Institut Agro, F-34398 Montpellier, France.

Published online: 02 June 2022

Reference

1. Su R, Dossou SSK, Dossa K, et al. Genome-wide characterization and identification of candidate *ERF* genes involved in various abiotic stress responses in sesame (*Sesamum indicum* L.). *BMC Plant Biol.* 2022;22:256. <https://doi.org/10.1186/s12870-022-03632-7>.

The original article can be found online at <https://doi.org/10.1186/s12870-022-03632-7>.

*Correspondence: wuzm@jxau.edu.cn; junyou@caas.cn

¹ Key Laboratory of Crop Physiology, Ecology, and Genetic Breeding, Ministry of Education/College of Agronomy, Jiangxi Agricultural University, Nanchang 330045, China

² Key Laboratory of Biology and Genetic Improvement of Oil Crops of the Ministry of Agriculture and Rural Affairs, Oil Crops Research Institute of the Chinese Academy of Agricultural Sciences, Wuhan 430062, China

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.