

Research article

Open Access

## Synthesis of L-ascorbic acid in the phloem

Robert D Hancock, Diane McRae, Sophie Haupt and Roberto Viola\*

Address: Unit of Plant Biochemistry, Scottish Crop Research Institute, Invergowrie, Dundee DD2 5DA, United Kingdom

Email: Robert D Hancock - rhanco@scri.sari.ac.uk; Diane McRae - dmcrae@scri.sari.ac.uk; Sophie Haupt - shaupt@scri.sari.ac.uk; Roberto Viola\* - rviola@scri.sari.ac.uk

\* Corresponding author

Published: 24 November 2003

Received: 03 September 2003

BMC Plant Biology 2003, 3:7

Accepted: 24 November 2003

This article is available from: <http://www.biomedcentral.com/1471-2229/3/7>

© 2003 Hancock et al; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

### Abstract

**Background:** Although plants are the main source of vitamin C in the human diet, we still have a limited understanding of how plants synthesise L-ascorbic acid (AsA) and what regulates its concentration in different plant tissues. In particular, the enormous variability in the vitamin C content of storage organs from different plants remains unexplained. Possible sources of AsA in plant storage organs include *in situ* synthesis and long-distance transport of AsA synthesised in other tissues via the phloem. In this paper we examine a third possibility, that of synthesis within the phloem.

**Results:** We provide evidence for the presence of AsA in the phloem sap of a wide range of crop species using aphid stylectomy and histochemical approaches. The activity of almost all the enzymes of the primary AsA biosynthetic pathway were detected in phloem-rich vascular exudates from *Cucurbita pepo* fruits and AsA biosynthesis was demonstrated in isolated phloem strands from *Apium graveolens* petioles incubated with a range of precursors (D-glucose, D-mannose, L-galactose and L-galactono-1,4-lactone). Phloem uptake of D-[U-<sup>14</sup>C]mannose and L-[1-<sup>14</sup>C]galactose (intermediates of the AsA biosynthetic pathway) as well as L-[1-<sup>14</sup>C]AsA and L-[1-<sup>14</sup>C]DHA, was observed in *Nicotiana benthamiana* leaf discs.

**Conclusions:** We present the novel finding that active AsA biosynthesis occurs in the phloem. This process must now be considered in the context of mechanisms implicated in whole plant AsA distribution. This work should provoke studies aimed at elucidation of the *in vivo* substrates for phloem AsA biosynthesis and its contribution to AsA accumulation in plant storage organs.

### Background

In plants, L-ascorbic acid (AsA) is essential for photosynthetic activity via the detoxification of superoxide and hydrogen peroxide in chloroplasts in the absence of catalase [1]. AsA is also crucially involved in the regeneration of  $\alpha$ -tocopherol and zeaxanthin and the pH-mediated modulation of PS II activity [2]. The critical importance of AsA in photosynthetic metabolism is emphasised by its abundance in chloroplasts where its concentration reaches 50 mM [2]. The constitutive role of AsA in photo-

synthesis explains its widespread distribution and generally high content in leaves [3,4]. These studies also revealed relatively low variability of AsA content between species, with coefficients of variation ranging between 59% in herbaceous plants (211 species) and 67% in woody plants (41 species). Conversely, the variability in the AsA content of non-photosynthetic tissues such as storage organs is generally much higher. Our estimates made from available data on fruits and vegetables from 65 species show an average vitamin C (AsA +

dehydroascorbic acid [DHA]) content of 1.1 mg gFW<sup>-1</sup> with a coefficient of variation of 334%. For example, a 10,000-fold difference in AsA content is found between the fruits of camu camu (*Miricaria dubia*), which contains up to 30 mg gFW<sup>-1</sup> vitamin C [5] and the medlar (*Mespilus germanica*) containing less than 3 µg gFW<sup>-1</sup> [6]. There is no obvious taxonomic explanation for such differences and large variations in storage organ AsA content are also observed within individual species including cultivated species such as strawberry and blackcurrant [7,8]. There is also no clear physiological explanation for the vast variability of AsA content in plant storage organs, but our understanding of AsA functions in these tissues is still very limited. In addition to its general antioxidant functions, AsA has been implicated in cell division, cell wall metabolism, cell expansion, and plant-pathogen interactions [9].

Advances in our understanding of AsA biosynthesis in plants have been made recently. There is general consensus that the biosynthetic pathway proposed by the Smirnoff group [10], represents a major pathway in plants. Confirmation has been obtained from analyses of *Arabidopsis thaliana* AsA deficient mutants [11,12] and from identification and manipulation of several of the genes involved [13-15]. However, there are reports of AsA accumulation in lettuce over-expressing rat gulono-lactone oxidase [16] and in *A. thaliana* constitutively expressing a strawberry D-galacturonic acid reductase [17] suggesting alternative pathways may be operational. Additionally, AsA accumulation has been reported in maize and tobacco constitutively over-expressing the AsA recycling gene DHA reductase [18]. Thus, despite this renewed interest, the mechanisms controlling AsA accumulation in storage organs remain largely elusive. It is still unclear for example whether AsA accumulation in storage organs occurs as a result of *in situ* synthesis or import from photosynthetically active regions. There is evidence for developmentally regulated AsA accumulation in fruits with high rates soon after anthesis [8] and in potato tubers a large accumulation is observed soon after tuber induction [19].

The competence for AsA synthesis seems to be ubiquitous amongst plant cells and AsA accumulation has been observed in a wide variety of cultured cells and excised tissues or organs incubated with appropriate substrates (e.g. [10,15] and [19-23]). However, evidence for direct AsA biosynthesis in storage organs *in planta* is missing. A recent report [24] described uptake of radiolabelled AsA by *Arabidopsis* and *Medicago* leaves and its movement to sink tissues such as shoot and root tips, flowers, siliques and sink leaves, leading the authors to suggest that long-distance transport of AsA occurs in plants. The presence of AsA in the phloem was first reported some time ago [25],

but its specific functions were not properly investigated. The purpose of the present paper is to provide additional information on phloem AsA content, specifically its origin and its implications for AsA accumulation in storage organs. By adopting a wide range of plant material and exploiting the most suitable model systems available we show that AsA is a widespread constituent of plant phloem sap, and that isolated phloem strands are competent for AsA biosynthesis from distant substrates. Moreover, we demonstrate that the intermediates of AsA biosynthesis D-mannose and L-galactose can be taken up by the phloem. The implications of these findings in relation to whole plant AsA distribution are discussed.


## Results

### Detection of AsA in plant vascular tissue

Aphid stylectomy was employed to isolate phloem exudates from a number of crop plants (barley, pea, potato, tobacco, turnip) in the absence of contamination from other tissues. When exudates were analysed by HPLC, AsA was detected in all cases (Fig. 1). We were unable to avoid evaporation of water from the exuded material and thus made no attempt to quantify AsA concentration in the exudates. However, it should be noted that in all cases AsA appeared as the dominant compound with absorption at 245 nm retained by the column.

In order to obtain more substantial amounts of exudate to allow quantification of AsA, we focussed our attention onto *Cucurbitaceae* fruits, from which relatively high volumes of vascular exudates (approximately 1 ml Kg<sup>-1</sup> of fruit) could be obtained. Table 1 shows the AsA concentration in vascular exudates from fruits of a number of *Cucurbitaceae* species in relation to that of soluble carbohydrates. Values in the region of 0.49 to 1.44 mg gFW<sup>-1</sup> (2.8 to 8.2 mM) were obtained, whilst the total carbohydrate concentration ranged from 10.95 to 23.28 mg gFW<sup>-1</sup>.

AsA localisation in the vascular tissue was confirmed by histochemical analysis of slices excised from courgette fruits and celery petiole (also used for metabolic studies, see below) using methanolic AgNO<sub>3</sub> solutions. This method exploits the specific ability of AsA to reduce Ag<sup>+</sup> at low temperature resulting in the formation of metallic silver deposits [26]. In both courgette fruit and celery petiole, metallic silver deposits were abundant in the vascular bundles with the staining in this tissue clearly limited to the phloem elements in the case of celery (Fig. 2). In addition, staining was evident in the tissue surrounding developing seeds in the courgette fruit and in the storage parenchyma in the celery petiole. No formation of metallic silver deposits were observed in control specimens pretreated with CuSO<sub>4</sub> to induce AsA oxidation. The penetration of AgNO<sub>3</sub> in parenchyma cells was confirmed by



**Figure 1**

**HPLC traces of plant phloem exudates obtained by aphid stylectomy.** Stylets were excised from aphids feeding on *Hordeum vulgare* (barley) (B), *Pisum sativum* (pea) (C), *Solanum tuberosum* (potato) (D), *Nicotiana tabacum* (tobacco) (E) and *Brassica rapa* ssp. *Oleifera* (turnip) (F) using a microcautery device. The exuded phloem sap was collected and diluted into ice-cold 5%  $H_3PO_3$  containing 5 mM TCEP. The figure shows chromatograms ( $A_{245}$ ) of phloem exudates and insets show the absorbance spectrum from the left and right sides of the peak with identical retention time to authentic AsA (A). The horizontal axis represents retention time (min) and the vertical axis detector response (milli-absorbance units).

**Table 1: Total carbohydrate and AsA content of *Curcubitaceae* fruit vascular exudates**

Species	Carbohydrate (mg gFW <sup>-1</sup> )	AsA (mg gFW <sup>-1</sup> )	AsA (mM)
<i>Cucumis sativus</i>	10.95 ± 0.13	1.02 ± 0.05	5.8 ± 0.33
<i>Cucurbita maxima</i>	22.48 ± 0.09	1.44 ± 0.04	8.2 ± 0.20
<i>C. moschata</i>	21.98 ± 0.80	0.49 ± 0.03	2.8 ± 0.15
<i>C. pepo</i> (courgette)	17.96 ± 0.34	0.87 ± 0.05	4.9 ± 0.26
<i>C. pepo</i> (squash)	20.46 ± 0.53	0.81 ± 0.02	4.6 ± 0.13
<i>C. maxima</i> x <i>C. moschata</i>	23.28 ± 0.66	0.63 ± 0.02	3.6 ± 0.09


Mean ± S.E. n = 3 Samples were extracted in 5%  $H_3PO_3$  (w/v) containing 5 mM TCEP and carbohydrate concentration estimated according to Dubois et al. (1956). AsA concentration was estimated by HPLC.

preincubation of tissues with 25 mM L-galactose which resulted in intense staining of parenchymal tissues (data not shown)

#### AsA biosynthesis in the phloem

The relatively large quantities of vascular exudates obtained from excision of courgette fruit enabled us to investigate the presence of the enzymes of the AsA biosyn-

thetic (Smirnoff-Wheeler) pathway (Fig. 3). We detected activities of all the previously identified enzymes of the AsA biosynthetic pathway [10] in the vascular exudates with the exception of the terminal (mitochondrial) enzyme L-galactono-1,4-lactone dehydrogenase (Table 2). Similarly, no conversion of L-galactono-1,4-lactone to AsA was detected in isolated vascular exudate (data not shown). Both vascular exudates and homogenised fruit




**Figure 2**

**Localisation of AsA with methanolic  $\text{AgNO}_3$  in courgette and celery tissues.** The left side of each image shows sections stained immediately after excision and the right side control sections which were pre-incubated in 5% (w/v)  $\text{CuSO}_4$  to oxidise endogenous AsA. Transverse section of courgette fruit (Ai) shows intense staining in the region of the outer vascular ring (arrows). Staining was also observed in the region surrounding developing seeds. Detail of the outer vascular area confirms that staining was confined to vascular regions (Aii). Staining was absent in control sections pre-incubated with 5%  $\text{CuSO}_4$ . In the celery petiole (Bi) intense staining was associated with vascular bundles (arrows) and storage parenchyma (tail-less arrows). Detail of the vascular region (Bii) shows intense staining associated with the phloem but not the xylem.

tissue were competent for the enzymic hydrolysis of GDP-L-[1- $^{14}\text{C}$ ]galactose although we were unable to test for the activity of L-galactose-1-phosphate phosphatase due to unavailability of substrate. In many cases, the enzyme activities in exudates were higher than in whole fruit tissue. In some cases e.g. phosphomannose isomerase, phosphomannose mutase and GDP-D-mannose pyrophosphorylase, activity was only detected in the vascular exudate. Incubation of L-[1- $^{14}\text{C}$ ]galactose directly in collected vascular exudate without any additions resulted in the formation of L-[ $^{14}\text{C}$ ]galactono-1,4-lactone (Fig. 4). This conversion was prevented by desalting of the exudate and could be restored by the addition of NAD to desalted exudate.

Functional characterisation of the AsA biosynthetic pathway in the phloem *in vivo* was carried out using isolated phloem strands from celery petioles, an established model system for phloem metabolism [27]. Phloem strands or storage parenchyma discs were incubated for 18 h with a range of AsA precursors. AsA accumulation was observed in both tissues upon incubation with L-galactose but the phloem strands accumulated ca. twice the amount of AsA compared with the parenchyma (Fig. 5). Incubation with L-galactono-1,4-lactone resulted in substantial AsA accumulation in the phloem strands only. AsA accumulation was not observed when samples were incubated with sucrose, D-glucose or D-mannose. However, the capacity of both tissues to synthesise AsA from these

**Figure 3**


**The Smirnov-Wheeler AsA biosynthetic pathway.** Plants synthesize AsA from glucose via phosphorylated hexose intermediates. 1. hexokinase (E.C. 2.7.1.1), 2. phosphoglucose isomerase (E.C. 5.3.1.9), 3. phosphomannose isomerase (E.C. 5.3.1.8), 4. phosphomannose mutase (E.C. 5.4.2.8), 5. GDP-mannose pyrophosphorylase (E.C. 2.7.7.22), 6. GDP-mannose-3,5-epimerase (E.C. 5.1.3.18), 7. GDP-L-galactose pyrophosphatase, 8. L-galactose-1-phosphate phosphatase, 9. L-galactose dehydrogenase, 10. L-galactono-1,4-lactone dehydrogenase (E.C. 1.3.2.3). Glc, D-glucose; Fru, D-fructose; Man, D-mannose; L-Gal, L-galactose; L-GalL, L-galactono-1,4-lactone. Where no E.C. numbers are listed they have yet to be assigned.

substrates was confirmed by incubation with D-[U-<sup>14</sup>C]glucose or D-[U-<sup>14</sup>C]mannose. Radiolabel accumulation into AsA was observed from both these substrates as well as from L-[1-<sup>14</sup>C]galactose (Table 3). With all labelled substrates, phloem strands showed a much higher total uptake and a significantly higher proportion of label incorporation into L-[<sup>14</sup>C]AsA compared with parenchyma discs.

#### **Uptake of AsA and other precursors by the leaf phloem**

The ability of source leaf phloem to take up L-[1-<sup>14</sup>C]AsA, or a number of substrates related to AsA metabolism ([1-<sup>14</sup>C]DHA, D-[U-<sup>14</sup>C]mannose, L-[1-<sup>14</sup>C]galactose and [U-

<sup>14</sup>C]sucrose as a reference) was tested in leaf discs of *Nicotiana benthamiana*, a model system for phloem uptake studies as it allows good resolution of major and minor veins [28]. Although [U-<sup>14</sup>C]sucrose and D-[U-<sup>14</sup>C]mannose appeared to be taken up more readily by the discs, the pattern of radioactivity distribution was similar with all precursors used, showing clear labelling of major and minor veins (Fig. 6). Except for L-[1-<sup>14</sup>C]ascorbic acid, no significant metabolism of applied substrates was observed during the incubation period. With L-[1-<sup>14</sup>C]ascorbic acid, ca. 80% of the radioactivity was present as [<sup>14</sup>C]DHA at the end of the experiment as determined by HPLC with flow scintillation analysis (data not shown).




**Figure 4**  
**Synthesis of L-[<sup>14</sup>C]galactono-1,4-lactone from L-[<sup>14</sup>C]galactose in courgette fruit exudate.** Radiochromatograms obtained from exudate incubated with L-[<sup>14</sup>C]galactose for 30 min at 30°C. The reaction was stopped by the addition of H<sub>3</sub>PO<sub>3</sub> prior to removal of precipitate and analysis by HPLC with online flow scintillation detection. Individual chromatograms show reactions containing L-[<sup>14</sup>C]galactose and heated (100 °C) exudate (A), fresh exudate (B), desalted exudate (C) and desalted exudate with addition of 0.5 mM NAD (D). The elution times of authentic L-[<sup>14</sup>C]galactose (L-Gal), L-[<sup>14</sup>C]galactono-1,4-lactone (L-GalL) and L-[<sup>14</sup>C]galactonic acid (L-GalA) are shown.

## Discussion

### Detection of AsA in the phloem

We have used stylectomy to access the phloem sap of a number of crop plants (barley, pea, potato, tobacco, turnip) and AsA was detected without exception. This finding corroborates earlier reports [24,25] and indicates that AsA is a widespread constituent of plant phloem. The AsA concentration in the samples obtained by stylectomy was not established in this or in previous studies but values of 0.49 to 1.45 mg gFW<sup>-1</sup> (2.8 to 8.2 mM) were obtained from vascular exudates of mature fruits of *Cucurbitaceae* species (Table 1). Such exudates have been used extensively to

study phloem metabolism [29-32] and have been shown to have properties typical of phloem saps such as high pH and inorganic ions, amino acid and organic acid levels similar to those of phloem saps from other species. The possibility of large dilution from xylem sap or water channels has also been dismissed [32] and the AsA concentration detected in vascular exudates from *Cucurbitaceae* species is comparable with the values reported by Ziegler [25] for other plant species. Additional evidence for the presence of AsA in the phloem tissue was obtained using a histochemical approach based on the ability of AsA to reduce cold, acidified, ethanolic AgNO<sub>3</sub> and generate



**Figure 5**


**Effect of exogenous substrates on AsA content of celery tissues.** Isolated phloem strands or storage parenchyma discs were incubated for 18 h in 20 mM MES pH 5.5, 300 mM mannitol, 5 mM MgCl<sub>2</sub>, 2 mM KCl, 1 mM CaCl<sub>2</sub>, 1 mM CaSO<sub>4</sub> (Daie, 1987) and 25 mM of the appropriate substrate. AsA was extracted and total AsA quantified by HPLC after reduction of DHA with 5 mM TCEP. Data represent mean ± S.E., n = 3.

metallic silver deposits [26]. In both courgette fruit and celery petioles, (i.e. the plant organs used for experimental investigation on AsA biosynthesis) strong staining was observed in the vascular region with metallic deposit specifically localised to the phloem tissue in the case of celery (Fig. 2). Although this technique is not specific for AsA, its selectivity is substantially improved at low temperature and pH where other reducing agents are ineffective [26,33]. Additionally, pre-treatment of plant specimens with CuSO<sub>4</sub> completely prevented the formation of metallic silver deposits (Fig. 2). The relatively more intense staining of the vascular region in relation to the surrounding parenchymous areas may be explained if AsA concentration in the cytosol was substantially higher than in the

vacuole [34]. Intense staining of non-vacuolated (meristematic) cells has been previously reported with this technique [35].

#### **Synthesis of AsA in the phloem**

The presence of AsA synthesising enzymes was investigated in vascular exudates from mature fruits of courgette, a system validated for the isolation and identification of phloem resident proteins [36]. With the exception of the putative L-galactose 1-phosphate phosphatase [10] and L-galactono-1,4-lactone dehydrogenase, the activities of all AsA biosynthetic enzymes were detected in the vascular exudates. This includes the enzyme activity responsible for the hydrolysis of GDP-L-galactose, a key rate-limiting

**Figure 6**

**Uptake of AsA related metabolites by *Nicotiana benthamiana* leaf discs.** *N. benthamiana* source leaf discs were prepared as described in Materials and Methods and incubated with 25 mM MES pH 5.5, 20 mM CaCl<sub>2</sub> and the appropriate labelled substrate (at a final S.A. of 18.5 kBq mmol<sup>-1</sup>) for 3 h in the light (200 μmol m<sup>-2</sup> s<sup>-1</sup>). Leaf discs were washed, dried and lyophilised prior to exposure to Kodak Biomax X-ray film for 1 week. With all substrates tested, labelling is observed in both the major and minor (arrows) leaf veins. Labelling intensity appeared much higher with sucrose and mannose compared with the other substrates.

**Table 2: Activity of Smirnoff Pathway Enzymes in Courgette Vascular Exudate and Tissue**

Activity <sup>a</sup>	Phloem	Tissue
HK (D-Glc)	23.0 ± 1.73	22.5 ± 2.89
HK (D-Frc)	29.7 ± 1.33	24.8 ± 3.58
HK (D-Man)	4.0 ± 0.40	4.0 ± 0.58
PGI	1861.3 ± 15.18	1194.7 ± 173.38
PMI	94.2 ± 5.20	N.D. <sup>b</sup>
PMM	1.5 ± 0.11	N.D. <sup>b</sup>
GDPM PPase	22.8 ± 1.56	N.D. <sup>b</sup>
GDPM-3,5-epimerase	0.07 ± 0.012	0.12 ± 0.017
GDP-L-gal PPPase	0.012 ± 0.002	0.010 ± 0.002
L-GalDH	28.0 ± 0.46	7.0 ± 1.27
L-GalLDH	N.D. <sup>b</sup>	1.98 ± 0.18

<sup>a</sup> Activity is expressed in nmol min<sup>-1</sup> gFW<sup>-1</sup> of tissue ± S.E., n = 3 <sup>b</sup>N.D. = Activity not detected Courgette fruit exudate or tissue was prepared and desalted as described in the text. Enzymes listed are: HK, hexokinase (E.C. 2.7.1.1) with D-glucose (D-Glc), D-fructose (D-Frc) or D-mannose (D-Man) as substrate; PGI, phosphoglucose isomerase (E.C. 5.3.1.9); PMI, phosphomannose isomerase (E.C. 5.3.1.8); PMM, phosphomannose mutase (E.C. 5.4.2.8); GDPM PPase, GDP-D-mannose pyrophosphorylase (E.C. 2.7.7.22); GDPM-3,5-epimerase, GDP-D-mannose-3,5-epimerase (E.C. 5.1.3.18); GDP-L-gal PPPase, GDP-L-galactose pyrophosphatase; L-GalDH, L-galactose dehydrogenase; L-GalLDH, L-galactono-1,4-lactone dehydrogenase (E.C. 1.3.2.3). Where no E.C. numbers are supplied they have yet to be assigned.

step for AsA biosynthesis in plants (Hancock and Viola, in preparation). To our knowledge this represents the first direct assay of GDP-L-galactose hydrolysis in extracts of

plant tissues. The enzyme activities in the exudates appeared substantial by comparison with whole tissue

**Table 3: Conversion of  $^{14}\text{C}$ -precursors to AsA in celery tissues**

Substrate Tissue	D-[U- $^{14}\text{C}$ ]glucose		D-[U- $^{14}\text{C}$ ]mannose		L-[1- $^{14}\text{C}$ ]galactose	
	Parenchyma	Vascular	Parenchyma	Vascular	Parenchyma	Vascular
% Incorporation to AsA <sup>a</sup>	1.4 ± 0.06	3.1 ± 0.23	2.7 ± 0.06	5.9 ± 0.23	19.9 ± 0.92	49.8 ± 0.46
Uptake ( $\mu\text{Ci gFW}^{-1}$ ) <sup>a</sup>	0.65 ± 0.07	3.61 ± 0.05	0.32 ± 0.02	3.49 ± 0.05	0.09 ± 0.01	0.67 ± 0.02
Uptake (nmol gFW <sup>-1</sup> ) <sup>a</sup>	2.1 ± 0.23	11.6 ± 0.17	1.0 ± 0.06	11.4 ± 0.17	1.6 ± 0.12	12.2 ± 0.23

<sup>a</sup> Values are presented as mean ± S.E., n = 3. Samples were prepared and incubated as described in the text. After incubation samples were extracted with 5% (w/v) PCA containing 5 mM TCEP and AsA purified as described in materials and methods. Radiolabel distribution to AsA was quantified by HPLC using online flow scintillation detection.

extracts and for some of the enzymes activity was detectable solely in the exudates.

However, this may again reflect the vacuolisation of parenchyma cells resulting in dilution of enzyme activities below our levels of detection. We also show that vascular exudates contain, in addition to the required enzyme activities, pyridine nucleotides required for the conversion of L-[1- $^{14}\text{C}$ ]galactose into L-[ $^{14}\text{C}$ ]galactono-1,4-lactone (Fig. 4). The further conversion of L-galactono-1,4-lactone into AsA would require the activity of L-galactono-1,4-lactone dehydrogenase, the only membrane-bound enzyme of the AsA biosynthetic pathway (located in the inner mitochondrial membrane [37]). It is generally acknowledged that mitochondria are not abundant in the phloem sap however it is expected that, *in vivo*, metabolites such as L-galactono-1,4-lactone may be exchanged between the sieve elements and the companion cells via specialised plasmodesmata [38]. Thus *in vivo* AsA biosynthesis within the phloem tissue (*i.e.* the sieve element-companion cell [SE-CC] complex) may involve enzymic cooperation between the sieve tubes and the mitochondria rich companion cells.

The competence of the phloem tissue for the *in vivo* conversion of L-galactose or L-galactono-1,4-lactone to AsA was demonstrated in isolated phloem strands from celery petioles (Fig. 5). Additionally incubation of the phloem strands with D-[U- $^{14}\text{C}$ ]glucose or D-[U- $^{14}\text{C}$ ]mannose resulted in substantial labelling of AsA clearly demonstrating that this tissue possesses a fully operational AsA biosynthetic (Smirnoff-Wheeler) pathway (Fig. 3). Moreover, AsA synthesis and partitioning to AsA from labelled precursors (including L-[1- $^{14}\text{C}$ ]galactose) was significantly higher in phloem strands compared with storage parenchyma tissue, although differences in rates of precursor uptake were observed between the tissues used. The overall conclusion from the different experimental approaches used is that the phloem unexpectedly represents a tissue with a highly active AsA biosynthetic capacity.

### Origin of phloem AsA

How can we reconcile our finding of phloem AsA biosynthesis with the hypothesis of long distance AsA transport in plants put forward by Franceschi and Tarlyn [24]? These authors reported that application of  $^{14}\text{C}$ -AsA to leaf flaps in *Medicago sativa* and *A. thaliana* resulted in its translocation to sink tissues. We also found that that supply of L-[1- $^{14}\text{C}$ ]AsA (and [1- $^{14}\text{C}$ ]DHA) to source leaf discs of *Nicotiana benthamiana* resulted in accumulation of radiolabel in both major and minor veins. However, care needs to be used in interpreting experiments where exogenous substrates are introduced into the plant transport system via abraded or damaged leaves. Evidence of phloem translocation (via mass flow) of xenobiotics supplied to leaves and their unloading in sinks is available from experiments with carboxyfluorescein diacetate and other phloem tracers [e.g. [39]]. Work with carboxyfluorescein diacetate also demonstrates that phloem leakage of charged molecules (e.g. AsA) during translocation to sink tissues is minimal. An alternative possibility is that phloem AsA biosynthesis may be directly relevant to long-distance transport of AsA. For example, the SE-CC is known to contain sucrose synthase which is involved in sucrolytic cleavage to sustain energy production [40]. Sucrolytic intermediates could thus be used to sustain AsA biosynthesis within the SE-CC complex via the Smirnoff-Wheeler pathway. Micro-tubers obtained from isolated potato internodes cultured *in vitro* in the dark with sucrose as the sole carbon source accumulated AsA in concentrations similar to those observed in tubers obtained from field-grown plants [19]. Potato tubers are phloem-rich organs with a network of phloem strands internal and external to the vascular ring [39] and AsA distribution in potato tubers closely follows the phloem distribution (Viola et al., unpublished). However, our finding that both D-[U- $^{14}\text{C}$ ]mannose and L-[1- $^{14}\text{C}$ ]galactose were readily taken up by *N. benthamiana* leaf discs (resulting in a labelling intensity of major and minor veins similar to that of [U- $^{14}\text{C}$ ]sucrose in the case of the former) also raises the possibility that more direct intermediates of AsA biosynthesis may be provided to the phloem by the leaf mesophyll. In a following paper we show that L-galactose

supply to source potato leaves results in AsA accumulation in developing tubers (Tedone et al., in preparation).

## Conclusions

We show here that AsA presence in the phloem of higher plants is widespread and that the SE-CC complex of courgette and celery is competent for the synthesis of AsA from a range of substrates. Further work is required to establish precisely the nature of the substrate for phloem AsA biosynthesis *in vivo* and to establish quantitatively the contribution of phloem biosynthesis to the overall AsA content in storage organs. Although AsA transporters will be required for the transfer of AsA into storage organs involving apoplastic unloading steps, phloem biosynthesis may represent a significant direct source of AsA for those storage organs (e.g. potato tubers) where phloem unloading follows a symplastic route [39].

## Methods

### Plant Material and Preparation

Celery (*Apium graveolens* L.) and courgette (*Curbita pepo* L.) were purchased locally. Other plants were grown in unheated glasshouses under conditions of natural light in a standard mixture of peat, sand, limestone, vermiculite and celcote containing sincrostart and sincrocel 6 fertilisers. Plants were hand watered daily. Celery vascular bundles and parenchyma discs were prepared as described [27]. Vascular exudates were collected from courgettes and other *Cucurbitaceae* by positive displacement pipette from mature cut fruit and directly transferred to the appropriate ice-cold buffer for either enzyme activity or AsA determinations (see below). Phloem exudates from other plants were collected directly from severed stylets of *Myzus persicae* feeding on source leaf petioles. Prior to HPLC injection, phloem exudate was diluted into ice-cold 5% metaphosphoric acid with or without 5 mM tris(2-carboxyethyl)phosphine hydrochloride (TCEP).

### Chemicals and labelled substrates

D-[U-<sup>14</sup>C]glucose (S.A. 11.47 MBq mmol<sup>-1</sup>), D-[U-<sup>14</sup>C]mannose (S.A. 11.36 MBq mmol<sup>-1</sup>), [U-<sup>14</sup>C]sucrose (S.A. 17.09 MBq mmol<sup>-1</sup>) and L-[1-<sup>14</sup>C]AsA (S.A. 0.52 MBq mmol<sup>-1</sup>) were obtained from Amersham Pharmacia Biotech, Little Chalfont, UK. L-[1-<sup>14</sup>C]galactose (S.A. 2.04 MBq mmol<sup>-1</sup>) was obtained from American Radiolabelled Chemicals, St. Louis, USA. [<sup>14</sup>C]DHA was obtained by incubating 0.037 MBq L-[1-<sup>14</sup>C]AsA with 10 µg ascorbate oxidase (E.C. 1.10.3.3; Roche, Lewes, UK) in 50 mM sodium phosphate buffer pH 5.6 (final volume 100 µl). L-[1-<sup>14</sup>C]galactonic acid was synthesised by incubation of 0.037 MBq L-[1-<sup>14</sup>C]galactose with 1U fucose dehydrogenase (E.C. 1.1.1.122; Sigma, Dorset, UK) and 0.5 mM NADP in 50 mM glycine-KOH buffer pH 9.5 (final volume 100 µl). L-[1-<sup>14</sup>C]galactonolactone was obtained by incubation of 0.037 MBq L-[1-<sup>14</sup>C]galactose with 1U

fucose dehydrogenase and 0.5 mM NADP in 50 mM MOPS buffer pH 7.8 (100 µl). GDP-L-[1-<sup>14</sup>C]galactose was a gift from Dr. Ken Lawrie, GaxoSmithKline Pharmaceuticals, Marlow, UK and was synthesised according to the method of Baisch and Ohrlein [41] from L-[1-<sup>14</sup>C]galactose [ARC, St. Louis, USA]. The final product had a specific activity 0.37 MBq mmol<sup>-1</sup>, radiochemical purity of >95% as estimated by anion exchange HPLC and overall purity >90% as estimated by <sup>1</sup>H and <sup>13</sup>C NMR.

With the exception of L-fucose dehydrogenase and phosphomannose isomerase, which were purchased from Sigma, Dorset, UK, all enzymes were purchased from Roche, East Sussex, UK.

### In Situ Staining of AsA using Acidic Silver Nitrate

Transverse sections (approximately 1 mm) of either celery petiole or courgette fruit were cut by hand and washed briefly in distilled water then stained and fixed as described by Chinoy [26]. Control reactions were undertaken in which AsA in the tissue was first oxidised by exposure to aqueous 5% CuSO<sub>4</sub> as described [26].

### Tissue Incubation, Metabolite Extraction and HPLC Analysis

Celery tissues were incubated in buffer A consisting of 20 mM MES pH 5.5, 300 mM mannitol, 5 mM MgCl<sub>2</sub>, 2 mM KCl, 1 mM CaCl<sub>2</sub> and 1 mM CaSO<sub>4</sub> [27] in petri dishes with rotary shaking of 100 rpm. For experiments with unlabelled precursors, the appropriate compound was added to a final concentration of 25 mM and the incubation continued for 18 h. For labelling experiments, 400 mg tissue were incubated in 650 µl buffer containing 111 kBq of either D-[U-<sup>14</sup>C]glucose, D-[U-<sup>14</sup>C]mannose or L-[1-<sup>14</sup>C]galactose in sealed vessels with 100 rpm rotary shaking for 4 h. For experiments with unlabelled precursors, AsA was extracted by grinding the celery tissue in liquid N<sub>2</sub> followed by the addition of 5 volumes of 5% (w/v) metaphosphoric acid and further homogenisation. Cell debris was removed by centrifugation (16000 g, 1°C, 5 min). The supernatant was used directly for analysis of reduced AsA or incubated with 5 mM TCEP (1 h, 4°C) prior to analysis for measurement of total AsA by HPLC [22]. Recoveries of authentic AsA added to tissue samples immediately prior to extraction exceeded 90%. In radiolabelling experiments the tissue was removed from the incubation medium and washed three times for 5 min in 5 ml of buffer A. The tissue was then blotted, ground in liquid N<sub>2</sub> and homogenised in 5 volumes of 5% perchloric acid (PCA) containing 5 mM TCEP. After standing in ice for 30 min, cell debris was removed by centrifugation (16000 g, 5 min, 1°C) and a pH indicator (BDH 4080 indicator, 20 µl ml<sup>-1</sup>) was added to the supernatant. The pH range of the sample was adjusted to 6–7 by the drop-wise addition of a solution of 5 M K<sub>2</sub>CO<sub>3</sub>. Insoluble KClO<sub>3</sub> salts were

removed by centrifugation (16000 g, 5 min, 1°C), the supernatant was divided into 500 µl aliquots and applied to SAX SPE cartridges (100 mg, acetate counter ion, Alltech, Carnforth, UK). After washing the column with 4 ml H<sub>2</sub>O, L-[<sup>14</sup>C]AsA was released with 4 ml 300 mM formic acid. After lyophilisation and resuspension in 200 µl H<sub>2</sub>O, L-[<sup>14</sup>C]AsA was further purified and quantified by HPLC as previously described [22] with the exception that radioactivity was detected and quantified using a Packard 150TR radiodetector in the homogenous mode (500 µl flow cell) with postcolumn addition of Ultima Flo M scintillant at 3 ml min<sup>-1</sup>. Under these conditions, recovery of authentic L-[<sup>14</sup>C]AsA added to the tissue samples immediately prior to extraction exceeded 85%. Total carbohydrate determination in *Cucurbitaceae* fruit exudates was carried out using the phenol-sulphuric acid method [42].

#### Extraction and Determination of Enzyme Activities

Hexose kinase (HK), phosphoglucose isomerase (PGI), phosphomannose isomerase (PMI), phosphomannose mutase (PMM), GDP-L-galactose pyrophosphatase (GDP-L-gal PPPase) and L-galactose dehydrogenase activities (L-GalDH) were extracted from courgette fruit tissue by grinding in a mortar and pestle in ice cold 50 mM HEPES pH 8.0, 5 mM MgCl<sub>2</sub>, 5 mM DTT, 1 mM EDTA, 1 mM EGTA, 1 mM benzamidine hydrochloride and 0.5 mM PMSF (1:3 w/v). Cell debris was removed by centrifugation (10000 g, 10 min, 1°C) and sample supernatants were desalted on Sephadex G25 PD10 columns (Amersham Biosciences, UK) equilibrated with the same buffer prior to use. GDP-D-mannose pyrophosphorylase activity was extracted and desalted as described above with the exception that the buffer used throughout consisted of 100 mM tris pH 7.5, 15 mM 2-mercaptoethanol and 0.5 mM PMSF as described by Keller *et al.* [13]. GDP-D-mannose-3,5-epimerase activity was extracted and desalted using 100 mM tris pH 7.6, 5 mM DTT and 1 mM EDTA [43].

L-Galactono-1,4-lactone dehydrogenase (L-GalLDH) activity was extracted as described [44] with the exception that mitochondria were collected by centrifugation at 30000 g. In all cases, vascular exudates were treated exactly as whole tissue with the exception that exudates were diluted directly into extraction buffer and were not ground in a mortar and pestle.

HK activity was determined in a reaction mixture consisting of enzyme extract, 50 mM HEPES pH 8.0, 5 mM MgCl<sub>2</sub>, 2 mM ATP, 0.5 mM NAD and 1 U NAD-dependent glucose-6-phosphate dehydrogenase (from *Leuconostoc mesenteroides*) in a final volume of 1 ml. The reaction was undertaken at 30°C and started by the addition of 2 mM D-glucose. Reaction kinetics were followed by measurement of NAD reduction at 340 nm in a Hitachi U-3010

spectrophotometer. Control reactions were undertaken using boiled extract. HK activity was also measured using fructose as substrate in which case the reaction mixture additionally contained 1 U PGI or mannose as substrate where the mixture contained 1 U PMI in addition to PGI.

PGI activity was determined in a reaction mixture consisting of enzyme extract, 50 mM HEPES pH 8.0, 5 mM MgCl<sub>2</sub>, 0.5 mM NAD, 5 µM ZnSO<sub>4</sub> and 1 U glucose-6-phosphate dehydrogenase in a final volume of 1 ml. The reaction was undertaken at 30°C and started by the addition of 2 mM D-fructose-6-phosphate. Reaction kinetics were measured as described for hexokinase. PMI activity was measured as for PGI but the reaction mixture additionally contained 1 U PGI and was started by the addition of D-mannose-6-phosphate. PMM activity was assayed by including 1 U PMI in addition to PGI and starting the reaction with D-mannose-1-phosphate. In all cases control reactions were undertaken using boiled enzyme preparations.

GDP-D-mannose pyrophosphorylase activity was measured as described [13] with the exception that 2 mM NaF was included in the reaction mixture and control reactions lacked inorganic pyrophosphate. The reaction was linear with time for up to 2 h. GDP-D-mannose-3,5-epimerase activity was determined as described [43]. GDP-L-galactose pyrophosphatase activity was determined in a reaction mixture consisting of enzyme extract, 50 mM HEPES pH 8.0 and 5 mM MgCl<sub>2</sub> in a volume of 0.5 ml. The reaction was started by the addition of 1.11 kBq GDP-L-[<sup>14</sup>C]galactose. The reaction was stopped by the addition of 500 µl 10% (w/v) activated charcoal followed by brief vortexing. The charcoal was removed by centrifugation (16000 g; 10 min; 1°C) and radioactivity in the supernatant estimated by scintillation counting (Packard Tri-Carb 2000CA).

L-GalDH activity was measured in a reaction mixture consisting of enzyme extract, 50 mM HEPES pH 8.0, 5 mM MgCl<sub>2</sub> and 0.5 mM NAD. The reaction was started by the addition of 2 mM L-galactose. Reaction conditions and kinetic measurements were as described for hexokinase. L-Galactose dehydrogenase activity was also analysed in crude fruit exudate by addition of L-[<sup>14</sup>C]galactose. Samples were incubated at 30°C and the reaction stopped by addition of exudate to an equal volume of 5% (w/v) H<sub>3</sub>PO<sub>3</sub>.

Precipitated material was removed by centrifugation and formation of L-[<sup>14</sup>C]galactonic acid monitored by HPLC on a Metacarb 87C 300 × 7.8 mm column (MetaChem Technologies Inc., Torrance, CA) maintained at 70°C with a mobile phase of ultrapure water flowing at 1 ml min<sup>-1</sup>. HPLC equipment used was as described in Hancock *et al.*

[22] with the exception that radioactivity was detected and quantified using a Packard 150TR radiodetector in the homogenous mode (500  $\mu$ l flow cell) with postcolumn addition of Ultima Flo M scintillant at 3 ml min<sup>-1</sup>.

L-GalLDH activity was assayed as described by Oba et al. [44] with the exception that the buffer additionally contained 0.03% triton X-100 and 0.1 mM KCN to prevent reoxidation of cytochrome C by cytochrome C oxidase [37]. In another deviation from the published protocol, the reaction was started by addition of L-galactono-1,4-lactone to 5 mM. The value for molar extinction coefficient of cytochrome c at 550 nm used was 25300.

### Source Leaf Loading and Autoradiography

Loading of source leaves with <sup>14</sup>C-assimilates was undertaken as described by Turgeon and Gowan [45]. Leaves were removed from turgid plants and moistened with buffer B (25 mM MES pH 5.5, 20 mM CaCl<sub>2</sub>). The abaxial surface was gently abraded with carborundum powder and 16 mm discs cut using a sharp cork borer. Leaf discs were floated on buffer B under light (200  $\mu$ mol m<sup>-2</sup> sec<sup>-1</sup>) for 15 h at 25 °C with gentle rotary shaking. Finally, discs were transferred to 10 ml fresh buffer B containing 1 mM [U-<sup>14</sup>C]sucrose, D-[U-<sup>14</sup>C]mannose, L-[1-<sup>14</sup>C]galactose, L-[1-<sup>14</sup>C]AsA or [1-<sup>14</sup>C]DHA at a final S.A. of 18.5 kBq mmol<sup>-1</sup>. Incubation was continued under light for a further 3 h and subsequently, leaf discs were removed and washed five times (20 ml, 5 min) in buffer B. Small samples of incubation buffer were taken for HPLC analysis to ensure no metabolism had occurred at the beginning and end of the incubation period. Discs were surface dried and lyophilised between filter paper held flat between metal plates secured with G-clamps. Dried leaves were exposed to Kodak Biomax single sided X-ray film for 1 week at -80 °C prior to development.

### Authors' contributions

RDH undertook the biochemical investigations assisted by DMR, was responsible for data preparation and assisted in writing the manuscript. SH undertook the histochemical staining, performed the uptake studies in *N. benthamiana* leaf discs and prepared the figures. RV conceived the study and drafted the manuscript.

### Acknowledgements

The authors wish to thank Dr. Ken Lawrie, GlaxoSmithKline Pharmaceuticals for the gift of GDP-L-[1-<sup>14</sup>C]galactose. The Scottish Executive Environment and Rural Affairs Department funded this work.

### References

- Noctor G, Foyer CH: **Ascorbate and glutathione: Keeping active oxygen under control.** *Annu Rev Plant Physiol Plant Mol Biol* 1998, **49**:249-279.
- Smirnoff N: **The function and metabolism of ascorbic acid in plants.** *Ann Bot* 1996, **78**:661-669.
- Jones E, Hughes RE: **Foliar ascorbic acid in some angiosperms.** *Phytochem* 1983, **22**:2493-2499.
- Jones E, Hughes RE: **A note on the ascorbic acid content of some trees and woody shrubs.** *Phytochem* 1984, **23**:2366-2367.
- Justi KC, Visentainer JV, de Souza NE, Matsushita M: **Nutritional composition and vitamin C stability in stored camu-camu (*Myrciaria dubia*) pulp.** *Archivos Latinoamericanos De Nutricion* 2000, **50**:405-408.
- Rodriguez MAR, Oderiz MLV, Hernandez JL, Lozano JS: **Determination of vitamin-C and organic-acids in various fruits by HPLC.** *J Chromatogr Sci* 1992, **30**:433-437.
- Sone K, Mochizuki T, Noguchi Y: **Variations in ascorbic acid content among strawberry cultivars and their harvest times.** *J Japan Soc Hort Sci* 1999, **68**:1007-1014.
- Viola R, Brennan RM, Davies HV, Sommerville L: **L-Ascorbic acid accumulation in berries of *Ribes nigrum* L.** *J Hort Sci Biotechnol* 2000, **75**:409-412.
- Smirnoff N, Wheeler GL: **Ascorbic acid in plants: Biosynthesis and function.** *Critical Rev Biochem Mol Biol* 2000, **35**:291-314.
- Wheeler GL, Jones MA, Smirnoff N: **The biosynthetic pathway of vitamin C in higher plants.** *Nature* 1998, **393**:365-369.
- Conklin PL, Norris SR, Wheeler GL, Williams EH, Smirnoff N, Last RL: **Genetic evidence for the role of GDP-mannose in plant ascorbic acid (vitamin C) biosynthesis.** *Proc Natl Acad Sci USA* 1999, **96**:4198-4203.
- Conklin PL, Saracco SA, Norris SR, Last RL: **Identification of ascorbic acid-deficient *Arabidopsis thaliana* mutants.** *Genetics* 2000, **154**:847-856.
- Keller R, Springer F, Renz A, Kossmann J: **Antisense inhibition of the GDP-mannose pyrophosphorylase reduces the ascorbate content in transgenic plants leading to developmental changes during senescence.** *Plant J* 1999, **19**:131-141.
- Tabata K, Oba K, Suzuki K, Esaka M: **Generation and properties of ascorbic acid-deficient transgenic tobacco cells expressing antisense RNA for L-galactono-1,4-lactone dehydrogenase.** *Plant J* 2001, **27**:139-148.
- Gatzek S, Wheeler GL, Smirnoff N: **Antisense suppression of L-galactose dehydrogenase in *Arabidopsis thaliana* provides evidence for its role in ascorbate synthesis and reveals light modulated L-galactose synthesis.** *Plant J* 2002, **30**:541-533.
- Jain AK, Nessler CL: **Metabolic engineering of an alternative pathway for ascorbic acid biosynthesis in plants.** *Mol Breeding* 2000, **6**:73-78.
- Agius F, Gonzalez-Lamothe R, Caballero JL, Munoz-Blanco J, Botella MA, Valpuesta V: **Engineering increased vitamin C levels in plants by overexpression of a D-galacturonic acid reductase.** *Nat Biotechnol* 2003, **21**:177-181.
- Chen Z, Young TE, Ling J, Chang S-C, Gallie DR: **Increasing vitamin C content of plants through enhanced recycling.** *Proc Natl Acad Sci USA* 2003, **100**:3525-3530.
- Viola R, Vreugdenhil D, Davies HV, Sommerville L: **Accumulation of L-ascorbic acid in tuberising stolon tips of potato (*Solanum tuberosum* L.).** *J Plant Physiol* 1998, **152**:58-63.
- Yoneyama T, Yasuda M, Sato S, Takebe M: **<sup>13</sup>CO<sub>2</sub> Feeding studies on the metabolism of carbohydrates, ascorbate and oxalate in spinach (*Spinacia oleracea* L.).** *Soil Sci Plant Nutr* 1997, **43**:1147-1151.
- Davey MW, Gilot C, Persiau G, Ostergaard J, Han Y, Bauw GC, van Montagu MC: **Ascorbate biosynthesis in *Arabidopsis* cell suspension culture.** *Plant Physiol* 1999, **121**:535-543.
- Hancock RD, Galpin JR, Viola R: **Biosynthesis of L-ascorbic acid (vitamin C) by *Saccharomyces cerevisiae*.** *FEMS Microbiol Lett* 2000, **186**:245-250.
- De Pinto MC, Tommasi F, De Gara L: **Enzymes of the ascorbate biosynthesis and ascorbate-glutathione cycle in cultured cells of tobacco bright yellow 2.** *Plant Physiol Biochem* 2000, **38**:541-550.
- Franceschi VR, Tarlyn NL: **L-Ascorbic acid is accumulated in source leaf phloem and transported to sink tissues in plants.** *Plant Physiol* 2002, **130**:649-656.
- Ziegler H: **Nature of transported substances.** In *Encyclopedia of Plant Physiology New Series Volume 1*. Edited by: Zimmermann MH, Milburn JA. Berlin: Springer-Verlag; 1975:59-100.
- Chinoy NJ: **On the specificity of alcoholic, acidic silver nitrate reagent for the histochemical localization of ascorbic acid.** *Histochemie* 1969, **20**:105-107.
- Daie J: **Sucrose uptake in isolated phloem of celery is a single saturable transport system.** *Planta* 1987, **171**:474-482.

28. Eschrich W, Fromm J: **Evidence for two pathways of phloem loading.** *Physiol Plant* 1994, **90**:699-707.
29. Richardson PT, Baker DA, Ho LC: **Assimilate transport in cucurbits.** *J Exp Bot* 1984, **35**:1575-1581.
30. Alosi MC, Melroy DL, Park RB: **The regulation of gelation of phloem exudate from Cucurbita fruit by dilution, glutathione, and glutathione-reductase.** *Plant Physiol* 1988, **86**:1089-1094.
31. Walz C, Juenger M, Schad M, Kehr J: **Evidence for the presence and activity of a complete antioxidant defence system in mature sieve tubes.** *Plant J* 2002, **31**:189-197.
32. Fiehn O: **Metabolic networks of Cucurbita maxima phloem.** *Phytochem* 2003, **62**:875-886.
33. Chinoy NJ, Sanjeevan AG: **On the specificity of alcoholic silver nitrate reagent for the histochemical localizatin of ascorbic acid. A reappraisal.** *Histochemistry* 1978, **56**:275-282.
34. Rautenkranz AAF, Li L, Mächler F, Märtinoia E, Oertli JJ: **Transport of ascorbic and dehydroascorbic acids across protoplast and vacuole membranes isolated from barley (Hordeum vulgare L. cv Gerbel) leaves.** *Plant Physiol* 1994, **106**:187-193.
35. Patel KR, Berlyn GP: **Chemical investigations on multiple bud formation in tissue cultures of Pinus coulteri.** *Can J Bot* 1983, **61**:575-585.
36. Haebel S, Kehr J: **Matrix-assisted laser desorption/ionization time of flight mass spectrometry peptide mass fingerprints and post source decay: a tool for the identification and analysis of phloem proteins from Cucurbita maxima Duch. separated by two-dimensional polyacrylamide gel electrophoresis.** *Planta* 2001, **213**:586-593.
37. Siendones E, Gonzalez-Reyes JA, Santos-Ocana C, Navas P, Cordoba F: **Biosynthesis of ascorbic acid in kidney bean. L-Galactono-γ-lactone dehydrogenase is an intrinsic protein located at the inner mitochondrial membrane.** *Plant Physiol* 1999, **120**:907-912.
38. Van Bel AJE, Ehlers K, Knoblauch M: **Sieve elements caught in the act.** *Trends Plant Sci* 2002, **7**:126-132.
39. Viola R, Roberts AG, Haupt S, Gazzani S, Hancock RD, Marmioli N, Machray GC, Oparka KJ: **Tuberization in potato involves a switch from apoplastic to symplastic phloem unloading.** *Plant Cell* 2001, **13**:385-398.
40. Geigenberger P, Langenberger S, Wilke I, Heineke D, Heldt HW, Stitt M: **Sucrose is metabolised by sucrose synthase and glycolysis within the phloem complex of Ricinus communis L. seedlings.** *Planta* 1993, **190**:446-453.
41. Baisch G, Ohrlein R: **Convenient chemoenzymatic synthesis of β-purine-diphosphate sugars (GDP-fucose-analogues).** *Bioorg Med Chem* 1997, **5**:383-391.
42. Dubois M, Gilles KA, Hamilton JK, Rebers PA, Smith F: **Colorimetric method for the determination of sugars and related substances.** *Anal Chem* 1956, **28**:350-356.
43. Wolucka BA, Davey MW, Boerjan W: **A high-performance liquid chromatography radio method for determination of L-ascorbic acid and guanosine 5'-diphosphate-L-galactose, key metabolites of the plant vitamin C pathway.** *Anal Biochem* 2001, **294**:161-168.
44. Oba K, Ishikawa S, Nishikawa M, Mizuno H, Yamamoto T: **Purification and properties of L-galactono-γ-lactone dehydrogenase, a key enzyme for ascorbic acid biosynthesis, from sweet potato roots.** *J Biochem* 1995, **117**:120-124.
45. Turgeon R, Gowan E: **Phloem loading in Coleus blumei in the absence of carrier-mediated uptake of export sugar from the apoplast.** *Plant Physiol* 1990, **94**:1244-1249.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

