

RESEARCH ARTICLE

Open Access



# A *Lotus japonicus* E3 ligase interacts with the Nod Factor Receptor 5 and positively regulates nodulation

Daniela Tsikou<sup>1,2</sup>, Estrella E. Ramirez<sup>2</sup>, Ioanna S. Psarrakou<sup>1</sup>, Jaslyn E. Wong<sup>2</sup>, Dorthe B. Jensen<sup>2</sup>, Erika Isono<sup>3</sup>, Simona Radutoiu<sup>2\*</sup> and Kalliope K. Papadopoulou<sup>1\*</sup> 

## Abstract

**Background:** Post-translational modification of receptor proteins is involved in activation and de-activation of signalling systems in plants. Both ubiquitination and deubiquitination have been implicated in plant interactions with pathogens and symbionts.

**Results:** Here we present *LjPUB13*, a PUB-ARMADILLO repeat E3 ligase that specifically ubiquitinates the kinase domain of the Nod Factor receptor NFR5 and has a direct role in nodule organogenesis events in *Lotus japonicus*. Phenotypic analyses of three LORE1 retroelement insertion plant lines revealed that *pub13* plants display delayed and reduced nodulation capacity and retarded growth. *LjPUB13* expression is spatially regulated during symbiosis with *Mesorhizobium loti*, with increased levels in young developing nodules.

**Conclusion:** *LjPUB13* is an E3 ligase with a positive regulatory role during the initial stages of nodulation in *L. japonicus*.

**Keywords:** E3 ligase, Nodulation, *Lotus japonicus*, PUB13, Ubiquitination, Symbiosis

## Background

The legume-rhizobia symbiosis leads to the formation of novel organs on the plant root, termed nodules. Rhizobia within nodule cells differentiate into bacteroids that fix atmospheric dinitrogen in exchange for plant carbohydrates. The symbiotic signalling process is initiated when rhizobia secrete nodulation (Nod) factors upon sensing flavonoids produced by compatible legumes. *Lotus japonicus* Nod factor receptors NFR1 and NFR5 and the corresponding proteins LYK3 and NFP in *Medicago truncatula* and, GmNFR1 and GmNFR5 in soybean [1–8] are crucial for perception of rhizobial Nod factors. Rhizobia enter into roots through infection threads (ITs) that, in most cases, initiate in epidermal root hair cells and progress to inner root tissues (reviewed in [9]). Formation of functional nodules requires two separate but tightly coordinated

developmental processes: bacterial infection and nodule organogenesis. Protein ubiquitination has been identified as being critical for these two signalling pathways.

Ubiquitination of proteins, by which proteins are tagged by ubiquitin and subsequently destined to be degraded by the proteasome, is a regulatory process essential for eukaryotic growth, development and response to interacting microbes [10]. In some cases, ubiquitination is important for regulating the activity or trafficking of the target protein [11]. Post-translational modification by ubiquitination is accomplished by a three-step process that involves ATP-dependent activation of ubiquitin by an E1 enzyme, followed by conjugation by an E2 enzyme and specific ubiquitin ligation to substrate proteins by direct interaction by an E3 ligase [12, 13].

E3 ligases are divided into families based on their mechanism of action and on the presence of specific E2 interacting domains, such as HECT, RING and U-box (reviewed in [14, 15]). The largest class of plant U-box (PUB) proteins is the ARMADILLO (ARM) domain-containing PUB proteins that contain tandemly-repeated ARM motifs located at the

\* Correspondence: [radutoiu@mbg.au.dk](mailto:radutoiu@mbg.au.dk); [kalpapad@bio.uth.gr](mailto:kalpapad@bio.uth.gr)

<sup>2</sup>Department of Molecular Biology and Genetics, Aarhus University, Gustav Wieds Vej, 8000 C Aarhus, Denmark

<sup>1</sup>Department of Biochemistry and Biotechnology, University of Thessaly, Biopolis, 41500 Larisa, Greece

Full list of author information is available at the end of the article



C-terminal [16, 17]. ARM repeat proteins are known to be involved in a number of different cellular processes including signal transduction, cytoskeletal regulation, nuclear import, transcriptional regulation, and ubiquitination. PUB-ARM proteins have been implicated in plant receptor-like kinase (RLK) signalling, with the ARM repeat domain mediating the binding of PUBs to the kinase domain [17, 18].

E3 ligases have been shown to play roles in the establishment of legume-rhizobium symbiosis (reviewed in [19]). *M. truncatula* LIN [20] and the orthologous gene from *L. japonicus*, CERBERUS [21], encode E3 ligases containing U-box, ARM and WD-40 repeats, and have been reported to control rhizobial infection inside root hairs. A second PUB-ARM E3 ubiquitin ligase in *M. truncatula*, MtPUB1, was identified as a negative regulator of nodulation by direct interaction with the receptor-like kinase LYK3 [22]. *M. truncatula* PUB1 is required for both rhizobial and arbuscular mycorrhiza (AM) endosymbiosis as it also directly interacts with the receptor kinase DMI2, a key component of the common symbiosis signalling pathway [23]. A member of the SEVEN IN ABSENTIA (SINA) family of E3 ligases, SINA4, was shown to interact with SYMRK receptor-like kinase in *L. japonicus* and be a negative regulator of rhizobial infection [24]. *LjnsRING*, a RING-H2 E3 ubiquitin ligase from *L. japonicus*, was also reported to be required for both rhizobial infection and nodule function [25]. Nevertheless, the mechanistic mode of action of the E3 ligases in the symbiotic interactions has not been elucidated, and the targets of all the symbiotic E3 ligases remain to be identified.

Here, we report the involvement of a PUB-ARM protein, *LjPUB13*, in the symbiotic interaction of *L. japonicus* with rhizobia. *LjPUB13* is phylogenetically a close relative of *Arabidopsis* PUB13, which has been implicated in plant responses to bacterial flagellin (flg22) [26] and, recently, fungal long-chain chitooligosaccharides (chitooctaoase) [27]. In *L. japonicus*, *LjPUB13* is involved in the establishment of a successful symbiosis, through the interaction of its ARM domain with the Nod factor receptor NFR5 and direct ubiquitination of the NFR5 kinase domain.


## Results

### *LjPUB13* encodes an active E3 ligase

A *Lotus japonicus* sequence encoding for a novel ARMADILLO repeat-containing protein was initially identified as the TC63883 in the DFCI Gene Index Database (<http://compbio.dfci.harvard.edu/tgi/tgipage.html>), while searching for ARMADILLO repeat proteins in *L. japonicus*. Initial search at transcript data in *Lotus* Base (<https://lotus.au.dk>) showed that this gene is expressed in nodules. The predicted protein has a typical Plant

U-box (PUB) E3 ubiquitin ligase sequence with a U-box motif followed by a C-terminal ARMADILLO (ARM) repeat domain (Fig. 1a). It encodes a 671 amino acid long protein with an estimated molecular mass of 73 kDa. Phylogenetic analysis showed that this protein is distinct from other previously characterised PUBs in legumes, but it is the closest homolog to *Arabidopsis thaliana* PUB13 (Additional file 1: Figure S1 and S2). We therefore designated the protein as *LjPUB13*. The encoded polypeptide shares 69% and 65% amino acid identity with *AtPUB13* (AT3G46510) and *AtPUB12* (AT2G28830), respectively. At the genomic level, this gene displays an exon-intron structure similar to *AtPUB13* [28], but different from *AtPUB12* ([www.arabidopsis.org](http://www.arabidopsis.org)), with four exons spanning a 4547 bp region on chromosome 3 (Fig. 1a).

The predicted E3 ligase enzymatic activity of *LjPUB13* was analysed by an in vitro ubiquitination assay,



performed with *LjPUB13* purified as a glutathione S-transferase (GST) fusion protein from *Escherichia coli*. Incubation of *LjPUB13* with E1 ubiquitin-activating and E2 ubiquitin-conjugated enzymes, ubiquitin and ATP resulted in *LjPUB13* polyubiquitination (Fig. 1b). Bands with a larger molecular weight than *LjPUB13*-GST and a protein ladder were detected by anti-GST and anti-ubiquitin antibodies, respectively, when *LjPUB13* was added to the reaction (Fig. 1b, lane 4). The observed E3 ligase activity was abolished in the absence of E1 or E2 enzymes (Fig. 1b, lanes 1 & 2). These results show that *L. japonicus* PUB13 is a functional E3 ligase, possessing auto-ubiquitination activity.


### The expression of *LjPUB13* is symbiotically regulated

To test if *LjPUB13* is regulated during symbiosis, we monitored its transcript levels in roots and nodules at 7, 14, 21 and 28 days post inoculation (dpi) with *Mesorhizobium loti*. No significant increase was detected in inoculated plants compared to uninoculated ones. Functional nodules, in general, had approximately 2 times less *LjPUB13* transcript compared to roots (Fig. 2a).

To complement the results from gene transcript analysis, we constructed a *LjPUB13<sub>pro</sub>::GUS* fusion and analysed the spatial and temporal regulation of *LjPUB13* promoter activity in *Agrobacterium rhizogenes* transformed roots expressing the transgene. In both uninoculated and inoculated roots, the *LjPUB13* promoter was active throughout the whole root, but strongest in the vascular bundle (Fig. 2b) and at the lateral root initiation sites (Fig. 2c). In *M. loti* inoculated plants, an increase in the promoter activity in the root zones, where nodule primordia formation occurs, was observed (Fig. 2d). Although strong *LjPUB13<sub>pro</sub>::GUS* expression was detected in young developing nodules at 7 dpi (Fig. 2d), in the fully developed mature nodules (28 dpi), *LjPUB13<sub>pro</sub>::GUS* expression was restrained to the nodule-root connection zone (Fig. 2f). Collectively, these results illustrate that *LjPUB13* transcript is present in symbiotically active root and young nodule tissues.


### *Ljpub13* mutants display growth defects and a delayed and reduced nodulation capacity

To investigate whether *LjPUB13* plays a role in the interaction between *L. japonicus* and the *M. loti* symbiont, we identified three mutant lines that possess LORE1 retroelement insertions [29, 30] in *LjPUB13* (Fig. 1a) and homozygous insertion lines were obtained for phenotypic analyses. The expression levels of *LjPUB13* in uninoculated plants of *pub13.1*, *pub13.2* and *pub13.3* mutant lines are 4-, 4.5- and 19-fold reduced, respectively, compared to



wild-type plants of the same age (Additional file 1: Figure S3).

We observed that in the absence of the *M. loti* symbiont, *L. japonicus pub13* mutants displayed a reduced growth phenotype; the root and shoot length are significantly shorter in all *pub13* mutants when compared to wild type plants of the same age (Fig. 3). In the presence of *M. loti*, *pub13* mutants formed a significantly lower number of nodules compared to the wild type Gifu, both in absolute number (Fig. 4a) and when normalized to root length (Fig. 4b). The latter indicates that the reduced nodulation phenotype does not correlate to the shorter root phenotype and is further supported by the direct comparison of *pub13.1* with the *har1* hypernodulation mutant [31, 32], where in *har1* very short roots (shorter than *pub13.1*) a high number of nodules is




formed (Additional file 1: Figure S4). Notably, nodulation was not only significantly reduced but also delayed in *pub13* mutants (Fig. 4c, Additional file 1: Figure S5). At 14 dpi only 15% of *pub13-1* and *pub13-2* plants had initiated symbiosis and had nodules compared to approximately 60% of wild type plants. This frequency in nodule appearance in the different plants of each plant line was higher at later stages of nodulation (Fig. 4c). Sections of mature nodules showed that they were fully infected and morphologically normal (Additional file 1: Figure S6). In addition, the root infection process was

not evidently perturbed in the *L. japonicus pub13* mutants since both wild type plants and *pub13* mutants formed a similar number of infection threads in root hairs (Fig. 4d). Together, these results show that *LjPUB13* is involved in plant growth and nodule organogenesis.

#### *L. japonicus pub13* mutants display normal root PTI responses to flg22 treatment

Based on its gene structure, amino acid identity and phylogenetic relationship, *LjPUB13* could be considered



a putative ortholog of *AtPUB13* in *L. japonicus*. In *Arabidopsis*, a number of phenotypes have been reported for *Atpub13* mutants, which primarily indicates for a role in PTI (pathogen-associated molecular patterns (PAMP)/ pattern-triggered immunity) response to flg22 treatment [26, 33]. Although the studies in *Arabidopsis* were focused in leaves, we searched for similar functions in *L. japonicus* roots as our aim was to investigate the function of *LjPUB13* in root responses to interacting microbes or microbial signals.

Firstly, we investigated the involvement of *LjPUB13* in the induction of reactive oxygen species (ROS) after flg22 treatment. Our analysis of wild type and *pub13* mutant roots revealed that similar levels of ROS were produced by wild type and mutant *L. japonicus* plants after treatment with 0.5  $\mu$ M flg22 (Additional file 1: Figure S7).

Secondly, we analysed the transcriptional activation of the defence marker genes *LjMPK3*, *LjPEROXIDASE* and *LjPR1*, together with *LjFLS2* and *LjBAK1* after 1-h treatment with flg22. All marker genes were induced in the treated samples (Additional file 1: Figure S8); *LjMPK3*, *LjPEROXIDASE* and *LjPR1* had a 4- to 6-fold increase in transcript levels while a 2- to 3-fold increase was observed for *LjFLS2* and *LjBAK1*. Nevertheless, the expression levels of these marker genes were comparable in wild type and *pub13* mutants in both treated and non-treated samples (Additional file 1: Figure S8), suggesting that *L. japonicus pub13* mutants neither express immunity-related genes constitutively nor overexpress them in the presence of flg22 as seen in *Arabidopsis* mutants [26].

Collectively, these results show that in *L. japonicus* roots the *PUB13* gene is not directly involved in PTI responses induced by flg22.


#### **Lotus PUB13 interacts with LjBAK1 but fails to ubiquitinate LjFLS2**

It is known that in *Arabidopsis*, PUB13, phosphorylated by BAK1 in the presence of flg22, polyubiquitinates the FLS2 receptor. This results in FLS2 degradation and regulation of the PTI signalling downstream of FLS2-BAK1 [26]. The apparent absence of PUB13-dependent PTI responses in *L. japonicus* roots prompted us to investigate the molecular basis of this differential response in *L. japonicus*.

We identified the corresponding *L. japonicus* BAK1, and the *LjBAK1<sub>cyt</sub>*-GST or *LjFLS2<sub>cyt</sub>*-GST fusion proteins were produced in *E. coli*. Next, we investigated the ability of the *LjPUB13*-HIS protein to interact with *LjBAK1* by performing an in vitro binding assay. Similar to *Arabidopsis* [26], we observed that *LjPUB13* interacts with *LjBAK1*, while the *LjPUB13*-*LjFLS2* interaction is barely detectable (Fig. 5a). *LjPUB13* appears to not be able to ubiquitinate *LjFLS2<sub>cyt</sub>* although we tested potential capacity in the presence of three different E2 enzymes (Additional file 1: Figure S9). These results indicate that PUB13 has different molecular capacities in *L. japonicus* and *Arabidopsis*.

#### **L. japonicus PUB13 interacts with and ubiquitinates NFR5**

Since *LjPUB13* is transcriptionally regulated during symbiosis and the gene is implicated in sustaining nodule organogenesis, we explored the involvement of *LjPUB13* in



**Fig. 5** *LjPUB13* interacts with *LjBAK1*, *LjFLS2* and *NFR5* in vitro. **a** In vitro binding assay of the HIS-tagged *LjPUB13* with the GST-fused cytoplasmic regions of *LjBAK1* and *LjFLS2*. *LjPUB13* interacts with *LjBAK1<sub>cyt</sub>* and *LjFLS2<sub>cyt</sub>*. **b** In vitro binding assays of the GST-fused *LjPUB13* and *LjBAK1<sub>cyt</sub>* with the HIS-tagged cytoplasmic regions of the Nod Factor receptors *NFR1* and *NFR5*. *LjPUB13* interacts strongly with *NFR5<sub>cyt</sub>*, while the *LjPUB13*-*NFR1<sub>cyt</sub>* interaction is undetectable. Sometimes a weak interaction of *LjPUB13*-*NFR1<sub>cyt</sub>* was observed but this result was not always reproducible. *LjBAK1<sub>cyt</sub>* also interacts with *NFR5<sub>cyt</sub>*, but no interaction was ever observed between *LjBAK1<sub>cyt</sub>* and *NFR1<sub>cyt</sub>*. After GST pulldown, bead-bound proteins were analyzed by immunoblotting using anti-HIS and anti-GST antibodies. These experiments were repeated five times

Nod factor signalling. Thus, we tested the possibility of *LjPUB13* to interact with the Nod factor receptors NFR1 and NFR5.

First, we investigated these supposed interactions *in vitro*. The cytoplasmic regions of NFR1 or NFR5 were expressed in *E. coli* as fusions to a HIS tag and were used in an *in vitro* binding assay together with the *LjPUB13*-GST fusion protein. We found that *LjPUB13* interacted strongly with NFR5<sub>cyt</sub> while the *LjPUB13*-NFR1<sub>cyt</sub> interaction was usually undetectable (Fig. 5b). Sometimes a weak interaction of *LjPUB13*-NFR1<sub>cyt</sub> was observed but this result was not always reproducible, and therefore cannot be fully considered as possible. Moreover, we tested whether *L. japonicus* BAK1, can interact with the receptors NFR1 and NFR5. *LjPUB13* strongly interacts with *LjBAK1* and it is plausible that *LjPUB13* acts together with *LjBAK1* as a complex. In addition, BAK1 is a well-known membrane co-receptor for many membrane receptor kinases [34]. Thus, in *in vitro* binding assays, GST-*LjBAK1*<sub>cyt</sub> was indeed found to interact with HIS-NFR5<sub>cyt</sub> but no interaction was observed between GST-*LjBAK1*<sub>cyt</sub> and HIS-NFR1<sub>cyt</sub> (Fig. 5b).


We verified the interaction of *LjPUB13* with the Nod factor receptors, using a BiFC (Bimolecular Fluorescence Complementation) assay in *Nicotiana benthamiana*. Since the ARM domain of PUB13 is responsible for the specificity in protein-protein interactions of E3 ligases, we created a truncated fusion protein, where only the ARM domain of *LjPUB13* (Additional file 1: Figure S2) was linked to N- or C-terminal half of YFP. We observed a strong reconstituted YFP signal on *N. benthamiana* leaf cells when *LjPUB13*<sub>ARM</sub> was co-expressed with *LjBAK1*, NFR1 or NFR5 (Fig. 6). Two other receptor-like kinases, *LjCLAVATA2* [35] and

*LjLYS11* [36], were used as negative control interactions. Indeed, no YFP signal was detected when *LjPUB13*<sub>ARM</sub> was co-expressed with either of the two control proteins. These results show that the ARM domain of *LjPUB13* recognizes and interacts with *L. japonicus* receptor kinases in a specific manner. In addition, the *LjBAK1*-NFR5 interaction was also confirmed by BiFC *in planta* (Fig. 6).


Finally, we examined whether *LjPUB13* can ubiquitinate the Nod factor receptors NFR1 and NFR5 since *LjPUB13* was shown to be a functional E3 ligase (Fig. 1b). The *E. coli*-produced HIS-NFR1<sub>cyt</sub> or HIS-NFR5<sub>cyt</sub> fusion proteins were used together with GST-*LjPUB13* in an *in vitro* ubiquitination assay. Interestingly, *LjPUB13* polyubiquitinated the cytosolic region of NFR5 and we detected high-molecular-weight proteins above the HIS-NFR5<sub>cyt</sub> (Fig. 7). The NFR5 ubiquitination by *LjPUB13* was demonstrated by using two different E2 enzymes, UbcH6 and UbcH5b. On the contrary, when HIS-NFR1<sub>cyt</sub> was tested as a substrate of *LjPUB13*, no ubiquitination activity was observed (Fig. 7). Thus, *L. japonicus* PUB13 specifically ubiquitinates the Nod factor receptor NFR5.

## Discussion

E3 ligases were identified as important signalling components in nitrogen-fixing symbiosis in model legumes [20–25]. However, the targets of all the symbiotic E3 ligases remain to be identified and the signalling events downstream of their action remain unexplored. Here, we report that a PUB-ARM E3 ligase that possesses ubiquitination activity (Fig. 1) plays a direct role in Nod factor signalling in *Lotus japonicus*, through its interaction and ubiquitination of the Nod factor receptor NFR5. *LjPUB13* is, thus, involved in the successful



**Fig. 6** *In planta* interactions of *LjPUB13*<sub>ARM</sub> and *LjBAK1* with the Nod Factor receptors NFR1 and NFR5. YFP split into N- and C-terminal halves were fused to the ARM domain of *LjPUB13* or to full length *LjBAK1*, NFR1 and NFR5. *N. benthamiana* leaves were cotransformed with different construct combinations and leaf epidermal cells were observed via confocal microscopy. The experiment was repeated four times with similar results. Bars, 50 μm



**Fig. 7** *LjPUB13* specifically ubiquitinates the Nod Factor receptor NFR5. *LjPUB13* was purified as GST fusion, while the cytoplasmic regions of NFR1 and NFR5 were purified as HIS fusion proteins. **a, b** The NFR5 ubiquitination was detected by an anti-HIS antibody, using two different E2 ligases: UbcH6 (**a**) and UbcH5b (**b**). No ubiquitination was observed when NFR1 was used as a substrate. **c** The presence of *LjPUB13* at the ubiquitination reactions was detected by an anti-GST antibody (input). **d** Purified NFR1-HIS and NFR5-HIS used at the ubiquitination reactions. The experiments were repeated three times with similar results

establishment of *L. japonicus*-rhizobium symbiosis, likely modulating the NFR5 protein levels or activity.

We found that *L. japonicus* PUB13 is involved in the organogenesis of nitrogen-fixing nodules. *pub13* mutant alleles displayed reduced and delayed nodule organogenetic capacity (Fig. 4). On the other hand, bacterial infection inside root hairs was normally sustained (Fig. 4d). This suggests that *LjPUB13* is required for the signalling events that lead to nodule organogenesis in the cortex, rather than the infection events. Furthermore, considering the low accumulation of *LjPUB13* gene transcripts in mature nodules (Fig. 2) and the successful colonization of nodules by rhizobium in the *pub13* mutants (Additional file 1: Figure S6), we envision that PUB13 E3 ligase plays a role in the initial stages of *L. japonicus*-rhizobium symbiosis establishment, rather than the later stages of nodule development and colonization.

The reduced nodulation phenotype could be attributed to the general reduced plant growth exhibited by *pub13* mutants (Fig. 3). However, the number of nodules was significantly different in *pub13* mutants from that of the wild-type plants also when expressed per unit of root length (Fig. 4b). This strongly suggests that a reduced nodulation is not directly related to a

shorter root. The fact that the nodulation capacity is independent of the shoot/root length is further supported by studies in hypernodulation mutants, like *har1* [31, 32], where an excessive number of nodules are produced by a very short plant (Additional file 1: Figure S4). Moreover, RNAi knockdown of *LjnsRING* E3 ligase, a nodule-specific gene involved in the early infection and mature nodule function [25], and mutation of *Amsh1*, a deubiquitination enzyme [37], affect plant growth in *L. japonicus*; this indicates that some proteins involved in (de)ubiquitination may act as nodes for plant growth and nodulation pathways. This is further supported by the expression of *LjPUB13* observed in developing lateral roots at non-inoculated plants (Fig. 2c). The growth defective phenotype of *L. japonicus pub13* mutants has also been reported for the *Arabidopsis pub13* mutants [38].

The expression of *LjPUB13* observed in nodule primordia (Fig. 2), together with the findings that *LjPUB13* interacts with (Figs. 5, 6) and directly ubiquitinates the NFR5 receptor (Fig. 7), strongly supports the requirement for *LjPUB13* during the early stages of nodule organogenesis in *L. japonicus*-rhizobia symbiosis. The defective nodulation phenotype observed in *pub13* mutants suggests a positive regulatory role of *LjPUB13* in

rhizobial symbiosis. On the contrary, the symbiotic E3s *MtPUB1* and *SINA4* [22, 24] exhibit a clear negative role on the early steps of the infection process. This indicates that different roles may be performed by various E3 ligases during symbiosis.

*LjPUB13* is the closest putative ortholog of *Arabidopsis PUB13* in *L. japonicus*. In *Arabidopsis*, *PUB13* has been implicated in FLS2-mediated flg22 signalling [26, 33] and, recently, in LYK5-mediated chitooctose responses [27]. We show here that the defence responses downstream of flg22 perception are independent of *PUB13* in the roots of this model legume. In contrast to what has been reported for *Arabidopsis* seedlings [26, 33], our results from ROS production, and defence gene regulation (Additional file 1: Figure S7 and S8) show that *LjPUB13* does not appear to be involved in flg22-dependent defence responses in *L. japonicus* roots. In line with this, our in vitro assays show that *LjPUB13* may not directly ubiquitinate *LjFLS2* (Additional file 1: Figure S9), as has been shown for its *Arabidopsis* counterpart [26]. However, we cannot rule out that the involvement of *LjPUB13* in plant immunity may be manifested in other parts of the plant or under conditions that have not been addressed in this study.

The differences observed in plant immunity responses in *L. japonicus pub13* mutant lines compared to *Arabidopsis* mutants, could be attributed to a host-dependent specialisation of function for these PUB proteins. Functional differentiation of orthologous PUB genes was also found for *Arabidopsis PUB17* and *Brassica napus ARC1* [39]. Alternatively, a redundancy in the roles of PUB proteins in *L. japonicus* could mean that paralogs may be responsible for the roles that have been assigned to *PUB13* in *Arabidopsis*. Blast analyses against *L. japonicus PUB13* in *Lotus* base v.3.0 revealed the presence of, yet uncharacterized, proteins with similarity to *PUB13*. Proteins with highest similarity (appr. 50% identity) are presented in Additional file 1: Figure S1. Future studies are needed to examine possible implication of these proteins in defence responses in *L. japonicus*.

Based on our results, we propose a plausible mechanism where *LjPUB13* acts on *NFR5* post-translationally. Ubiquitination of *NFR5* may lead to degradation, modulation of activity or re-localization. In any case, it is expected that *NFR5* turnover is essential for efficient nodule organogenesis, and the recruitment of *LjPUB13* ensures the onset and/or continuation of *NFR5*-mediated signalling and the successful initiation of nodule formation.

Along this line, and although *NFR5* internalization from the plasma membrane has not been shown directly as yet, an association of *NFR5* with the clathrin-mediated endocytosis has been proposed [40]. A clathrin protein (*CHC1*) was shown to interact with the Rho-like GTPase *ROP6* [41], an interaction partner of *NFR5* [42] in *L. japonicus*.

Interestingly, *ROP6* silencing in roots by RNAi did not affect the rhizobium entry in root hairs, but inhibited the IT growth through the root cortex, which resulted in the development of fewer nodules per plant [42].

We also show that *LjPUB13* physically and specifically interacts with the cytosolic regions of *LjBAK1* and *NFR5* in vitro and in vivo. Interestingly, the cytosolic region of *LjBAK1* was also shown to interact with the cytosolic region of *NFR5* (Figs. 5 and 6), suggesting that *PUB13/BAK1* may act as a complex in the establishment of *L. japonicus*-rhizobium symbiosis. The recruitment of *LjBAK1* in this case is anticipated, considering the ubiquitous role of *BAK1* in interactions of multiple membrane receptors kinases [26, 34].

## Conclusions

Ubiquitination and deubiquitination of proteins play a major role in root nodule symbiosis [20–25, 37]. Although many symbiotic E3 ligases have been identified to date, the mechanistic mode of action in the symbiotic interactions has not been elucidated, and their targets remain to be identified.

In the present study, the *L. japonicus* PUB-ARM E3 ligase *PUB13* was found to directly ubiquitinate the Nod factor receptor *NFR5*, and the delayed nodulation phenotype of the *pub13* mutants reveal the implication of this gene in nodule development. Based on our results, we suggest that *LjPUB13* has a role in the establishment of the *L. japonicus*-rhizobium symbiosis and acts as a positive regulator in nodule formation through the post-transcriptional control of *NFR5*.

## Methods

### Biological material, growth conditions and inoculation

*L. japonicus* ecotype Gifu B-129 was used as a wild type control. Wild type, *pub13* and *har1–3* [31] seeds were surface-sterilized as previously described [43] and were grown on wet filter paper for 3 to 4 days. Plants were then grown in petri dishes with solid quarter-strength B&D medium [44] on filter paper. The plants were grown in a vertical position in growth boxes, keeping the roots in the dark. Growth chamber conditions were 16-h day and 8-h night cycles at 21 °C.

For nodulation kinetics and IT counting, each petri dish was inoculated with 500 µl of a 0.02 OD<sub>600</sub> culture of *Mesorhizobium loti* cv. R7A DsRed [45]. Nodule numbers were scored on at least 30 plants and ITs were counted on at least 30 cm of root.

For gene expression analyses the plants were grown in pots with 2:1 mix of sand and vermiculite. Half of the plants were inoculated with a 0.1 OD<sub>600</sub> culture of *M. loti* cv. R7A. The plants were watered periodically with Hoagland nutrient solution.



### Protein purification

For GST-*LjPUB13*, GST-*LjBAK1<sub>cyt</sub>* and GST-*LjFLS2<sub>cyt</sub>*, the corresponding ORFs were PCR amplified and the resulting fragments were cloned between the EcoRI and XhoI sites of pGEX-6P-1 (GE Healthcare). For HIS-*LjPUB13*, the ORF was PCR amplified and the resulting fragment was cloned between the EcoRI and XhoI sites of pET21a (Novagen). Primers are listed in Additional file 1: Table S1. GST- and HIS- tagged proteins were expressed in *E. coli* Rosetta (DE3) (Merck Chemicals). GST-tagged proteins were then purified with Glutathione-Sepharose 4B beads (GE Healthcare) and eluted with 0.2 M reduced glutathione, whilst HIS-tagged proteins were purified with Ni Sepharose High Performance beads (GE Healthcare) and eluted with 250 mM imidazole.

Plasmids with the *NFR1<sub>cyt</sub>* and *NFR5<sub>cyt</sub>* fragments cloned in pProEX-1 vector were kindly provided by Mickael Blaise. HIS-*NFR1<sub>cyt</sub>* and HIS-*NFR5<sub>cyt</sub>* proteins were expressed in Rosetta 2 *E. coli* (DE3) competent cells (Novagen). IMAC purification was performed using Ni-NTA columns (Qiagen). The proteins were eluted with an elution buffer containing 50 mM Tris-HCl pH 8, 500 mM NaCl, 500 mM imidazole pH 8, 1 mM benzamidine, 5 mM  $\beta$ -mercaptoethanol and 10% glycerol. The eluted proteins were then injected onto a Superdex 200 increase 10/300 GL (GE Healthcare) column connected to an ÄKTA PURIFIER system (GE Healthcare) and eluted with a buffer containing 50 mM Tris-HCl pH 8, 500 mM NaCl, 5 mM  $\beta$ -mercaptoethanol and 10% glycerol.

### In vitro ubiquitination assay

The in vitro ubiquitination assays were performed with an Ubiquitylation kit (BML-UW9920, Enzo Life Sciences), using either UbcH6 or UbcH5b E2 enzymes and following the manufacturer's protocol. For *LjFLS2* ubiquitination tests UbcH5c E2 enzyme was also used.

All the reactions were incubated at 30 °C for 3 h, and then stopped by adding SDS sample buffer and boiled at 98 °C for 5 min. The samples were then separated by SDS-PAGE and analyzed by Western blotting, using the a-GST antibody (GE Healthcare) and the a-HIS antibody (Roche) to detect the tagged proteins or the anti-ubiquitin antibody (Santa Cruz Biotechnology P4D1) to detect the ubiquitinated fraction.

### In vitro binding assay

For the in vitro binding assays, 2  $\mu$ g of a GST tagged protein was incubated with 2  $\mu$ g of a HIS tagged protein, in different protein combinations, in 150  $\mu$ l of cold buffer A (50 mM Tris-HCl pH 7.5, 100 mM NaCl, 10% glycerol) with 0.1% Triton X-100, for 1.5 h at 4 °C. Glutathione-Sepharose 4B beads (GE Healthcare) were added to the mixtures and incubated with the proteins

for 2 more hrs at 4 °C. The beads were then washed 3 times with cold buffer A. The bead-bound proteins were analysed by immunoblotting, according to standard protocols, using anti-GST (GE Healthcare) and anti-HIS (Roche) antibodies.

### Expression analysis by qRT-PCR

To test the temporal and spatial expression of *LjPUB13* in non-inoculated and *M. loti* inoculated *L. japonicus* plants and the expression of defence genes after treatment with flg22, analysis by qRT-PCR was performed as previously described [45, 46]. Gene primers are listed in Additional file 1: Table S2.

### LjPUB13 promoter activity in *L. japonicus*

For the promoter-GUS-terminator construction, a 1475 bp promoter with a 5' untranslated region (UTR) and a 310 bp terminator region was amplified from *L. japonicus* genomic DNA (the primers are listed in Additional file 1: Table S3) and cloned into GoldenGate vectors [47]. The PCR fragments were firstly cloned into GoldenGate Level 0 vectors before being assembled as a construct (promoter:GUS:terminator) in a modified pIV10 vector [48]. The construct was transferred into *A. rhizogenes* AGL1.

Hairy root induction using *A. rhizogenes* was performed as described previously [49]. Chimeric plants were transferred into magenta growth boxes containing a sterilized 4:1 mix of clay granules and vermiculite as well as ¼ strength B&D medium supplemented with 1 mM KNO<sub>3</sub>. For inoculation, liquid cultures of *M. loti* cv. R7A expressing DsRed were grown to an optical density of 0.02 and applied directly to the root systems (0.7 ml per plant). Plants were grown at 21 °C (16 h light, 8 h dark) and harvested at indicated times post inoculation. GUS staining was performed as described previously [50]. Whole roots were visualized on a Leica M165 FC stereomicroscope.

### Infection thread formation

For inspection of infection-thread formation, roots were harvested 10 and 14 days after inoculation with *M. loti* strain cv. R7A DsRed. Sections (1 cm) of at least 30 plants were examined under a Zeiss Axioplan 2 fluorescent microscope.

### ROS accumulation

Seven-day-old roots were cut into 0.5 cm pieces and incubated overnight (with shaking) in 200  $\mu$ l water in 96-well plates (Grenier Bio-one). Before the measurements, the water was exchanged with 200  $\mu$ l of buffer (20 mM luminol, Sigma; 5  $\mu$ g/ml horseradish peroxidase, Sigma), supplemented with either H<sub>2</sub>O or 0.5  $\mu$ M flg22 peptide (FLS22-P-1, Alpha Diagnostic). Luminescence

was recorded with a Varioskan™ Flash Multimode Reader (Thermo).

### Protein-protein interaction studies by bimolecular fluorescence complementation (BiFC)

The ARM domain of *LjPUB13* (Additional file 1: Figure S2) and the full length *LjBAK1* were cloned, using Gateway technology (Invitrogen), into the pGREEN029:35S:GW:-nYFP/cYFP vectors creating N- and C-terminal fusions to YFP. The primers are listed in Additional file 1: Table S3. The NFR1 and NFR5 fused to nYFP/cYFP constructs used were the same as those described in [51].

*Agrobacterium tumefaciens* AGL1 cells transformed with the protein expression plasmids were grown in 5 ml LB medium supplemented with appropriate antibiotics at 28 °C. Bacteria were pelleted by centrifugation at 4000 g for 20 min at room temperature and re-suspended in agroinfiltration medium (10 mM MgCl<sub>2</sub>, 10 mM MES and 450 μM acetosyringone), incubated for 2–3 h in the dark and finally re-suspended to an OD<sub>600</sub> of 0.2. A 1:1 mixture of cultures was prepared for each construct combination together with the P19 construct (to an OD<sub>600</sub> of 0.02). A 1-ml syringe was used for infiltration of the bacterial mixture to the abaxial side of the *Nicotiana benthamiana* leaves. Fluorescence was detected using a Zeiss LSM510 confocal microscope.

### Statistical analyses

Differences in the tested biological parameters between mutant and wild type plants were analyzed by Student's t-test. A significant level of 5% was applied.

### Additional file

**Additional file 1: Table S1.** Primers used for cloning into expression vectors. **Table S2.** Primers used in qRT-PCR. **Table S3.** Primers used for cloning into GoldenGate and Gateway vectors. **Figure S1.** Phylogenetic tree of amino acid sequences of *LjPUB13* with previously characterized E3s of other species and *L. japonicus* uncharacterized PUBs. **Figure S2.** Amino acid sequence alignment of *LjPUB13* with *AtPUB13*. **Figure S3.** Expression levels of *LjPUB13* in *pub13* LORE1 mutants. **Figure S4.** Formation of nodules in *pub13.1* vs. *har1* mutants. **Figure S5.** Inoculated 28-day-old wild type and homozygous *pub13.3* mutants. **Figure S6.** Nodule sections of wild type, *pub13.1* and *pub13.2* plants. **Figure S7.** ROS accumulation in roots of *pub13* mutants vs wt plants. **Figure S8.** Expression of defence genes in *L. japonicus* wild type and *pub13* mutants. **Figure S9.** *LjFLS2* ubiquitination tests. (DOCX 39900 kb)

### Abbreviations

ARM: Armadillo; BiFC: Bimolecular Fluorescence Complementation; cyt: Cytoplasmic; GST: Glutathione S-transferase; HIS: Poly-histidine; PTI: Pathogen-associated molecular patterns (PAMP)/pattern-triggered immunity; PUB: Plant U-box; ROS: Reactive oxygen species; YFP: Yellow Fluorescent Protein

### Acknowledgements

The authors wish to thank Prof. Jens Stougaard (Aarhus University) for scientific support and for making corrections and comments on the manuscript; and Prof. Claus Schwechheimer (Technische Universität

München) for hosting DT in his laboratory by providing financial support through the SFB924 and for making comments on the manuscript. The authors thank Mickael Blaise for providing the plasmids for HIS-NFR1<sub>cyt</sub> and HIS-NFR5<sub>cyt</sub> protein expression, Zoltán Bozsóki for helping with the ROS assay, and Finn Pedersen for taking care of the plants in the greenhouse.

### Funding

This work was partially supported by the Postgraduate Programs 3817 & 3439 of the Department of Biochemistry and Biotechnology, University of Thessaly (to KKP). DT and SR were supported by the Danish National Research Foundation grant no. DNR79. DT was supported by a STSM (290713–031130) from Cost Action FA1103.

### Availability of data and materials

All data generated or analysed during this study are included in this published article and its supplementary information files.

The *LjPUB13* sequence is available in GenBank under the accession number KY131979 and in *Lotus* base v.3.0 (<https://lotus.au.dk/>) as Lj3g3v3189730.1. Other sequence data from this article can be found in the GenBank under the following accession numbers: *LjBAK1* (KY131980), *LjFLS2* (JN099749), *NFR1* (AJ575249), *NFR5* (AJ575255).

### Authors' contributions

KKP, SR and DT conceived the research plan; DT, EER, ISP, JEW and DBJ conducted experiments; KKP, SR and EI supervised the experiments; EI provided training to DT in biochemistry; DT, SR and KKP analysed data; DT wrote the article with contributions of all the authors; SR and KKP complemented and revised the writing. All authors read and approved the final manuscript.

### Ethics approval and consent to participate

Not applicable.

### Consent for publication

Not applicable.

### Competing interests

The authors declare that they have no competing interests

### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

### Author details

<sup>1</sup>Department of Biochemistry and Biotechnology, University of Thessaly, Biopolis, 41500 Larisa, Greece. <sup>2</sup>Department of Molecular Biology and Genetics, Aarhus University, Gustav Wieds Vej, 8000 C Aarhus, Denmark. <sup>3</sup>Department of Plant Systems Biology, Technical University of Munich, Emil-Ramann-Strabe 4, Freising, Germany.

Received: 17 March 2018 Accepted: 13 September 2018

Published online: 03 October 2018

### References

- Limpens E, Franken C, Smit P, Willemsse J, Bisseling T, Geurts R. LysM domain receptor kinases regulating rhizobial nod factor-induced infection. *Science*. 2003;302:630–3.
- Madsen EB, Madsen LH, Radutoiu S, Olbryt M, Rakwalska M, Szczyglowski K, Sato S, Kaneko T, Tabata S, Sandal N, Stougaard J. A receptor kinase gene of the LysM type is involved in legume perception of rhizobial signals. *Nature*. 2003;425:637–40.
- Radutoiu S, Madsen LH, Madsen EB, Felle HH, Umehara Y, Grönlund M, Sato S, Nakamura Y, Tabata S, Sandal N, Stougaard J. Plant recognition of symbiotic bacteria requires two LysM receptor-like kinases. *Nature*. 2003;425: 585–92.
- Arrighi JF, Barre A, Ben Amor B, Bersoult A, Soriano LC, Mirabella R, de Carvalho-Niebel F, Journet EP, Ghéradin M, Huguet T, Geurts R, Dénarié J, Rougé P, Gough C. The *Medicago truncatula* lysine motif receptor-like kinase gene family includes NFP and new nodule-expressed genes. *Plant Physiol* 2006;142:265–279.

5. Mulder L, Lefebvre B, Cullimore J, Imbert A. LysM domains of *Medicago truncatula* NFP protein involved in nod factor perception. Glycosylation state, molecular modeling and docking of chitoooligosaccharides and nod factors. *Glycobiology*. 2006;16:801–9.
6. Smit P, Limpens E, Geurts R, Fedorova E, Dolgikh E, Gough C, Bisseling T. *Medicago* LYK3, an entry receptor in rhizobial nodulation factor signalling. *Plant Physiol*. 2007;145:183–91.
7. Indrasumunar A, Kereszt A, Searle I, Miyagi M, Li D, Nguyen CD, Men A, Carroll BJ, Gresshoff PM. Inactivation of duplicated nod factor receptor 5 (NFR5) genes in recessive loss-of-function non-nodulation mutants of allotetraploid soybean (*Glycine max* L. Merr.). *Plant Cell Physiol*. 2010;51:201–14.
8. Indrasumunar A, Searle I, Lin MH, Kereszt A, Men A, Carroll BJ, Gresshoff PM. Nodulation factor receptor kinase 1 $\alpha$  controls nodule organ number in soybean (*Glycine max* L. Merr.). *Plant J*. 2011;65:39–50.
9. Oldroyd GE, Murray JD, Poole PS, Downie JA. The rules of engagement in the legume-rhizobial symbiosis. *Annu Rev Genet*. 2011;45:119–44.
10. Vierstra RD. The ubiquitin–26S proteasome system at the nexus of plant biology. *Nat Rev Mol Cell Biol*. 2009;10:385–97.
11. Komander D. The emerging complexity of protein ubiquitination. *Biochem Soc Trans*. 2009;37:937–53.
12. Hershko A, Ciechanover A. The ubiquitin system. *Annu Rev Biochem*. 1998;67:425–79.
13. Pickart CM, Eddins MJ. Ubiquitin, structures, functions, mechanisms. *Biochim Biophys Acta*. 2004;1695:55–72.
14. Smalle J, Vierstra RD. The ubiquitin 26S proteasome proteolytic pathway. *Annu Rev Plant Biol*. 2004;55:555–90.
15. Stone SL, Callis J. Ubiquitin ligases mediate growth and development by promoting protein death. *Curr Opin Plant Biol*. 2007;10:624–32.
16. Mudgil Y, Shiu SH, Stone SL, Salt JN, Goring DR. A large complement of the predicted *Arabidopsis* ARM repeat proteins are members of the U-box E3 ubiquitin ligase family. *Plant Physiol*. 2004;134:59–66.
17. Samuel MA, Salt JN, Shiu SH, Goring DR. Multifunctional arm repeat domains in plants. *Int Rev Cell Mol Biol*. 2006;253:1–26.
18. Gu T, Mazzurco M, Sulaman W, Matias DD, Goring DR. Binding of an arm repeat protein to the kinase domain of the S-locus receptor kinase. *Proc Natl Acad Sci U S A*. 1998;95:382–7.
19. Hervé C, Lefebvre B, Cullimore J. How many E3 ubiquitin ligase are involved in the regulation of nodulation? *Plant Signal Behav*. 2011;6:660–4.
20. Kiss E, Oláh B, Kaló P, Morales M, Heckmann AB, Borbola A, Lóza A, Kontár K, Middleton P, Downie JA, et al. LIN, a novel type of U-box/WD40 protein, controls early infection by rhizobia in legumes. *Plant Physiol*. 2009;151:1239–49.
21. Yano K, Shibata S, Chen WL, Sato S, Kaneko T, Jurkiewicz A, Sandal N, Banba M, Imaizumi-Anraku H, Kojima T, et al. CERBERUS, a novel U-box protein containing WD-40 repeats, is required for formation of the infection thread and nodule development in the legume-rhizobium symbiosis. *Plant J*. 2009;60:168–80.
22. Mbengue M, Camut S, de Carvalho-Niebel F, Deslandes L, Froidure S, Klaus-Heisen D, Moreau S, Rivas S, Timmers T, Hervé C, et al. The *Medicago truncatula* E3 ubiquitin ligase PUB1 interacts with the LYK3 symbiotic receptor and negatively regulates infection and nodulation. *Plant Cell*. 2010;22:3474–88.
23. Vernié T, Camut S, Camps C, Rembliere C, de Carvalho-Niebel F, Mbengue M, Timmers T, Gascioli V, Thompson R, le Signor C, Lefebvre B, Cullimore J, Hervé C. PUB1 interacts with the receptor kinase DMI2 and negatively regulates rhizobial and arbuscular mycorrhizal symbioses through its ubiquitination activity in *Medicago truncatula*. *Plant Physiol*. 2016;170:2312–24.
24. Den Herder G, Yoshida S, Antolín-Llovera M, Ried MK, Parniske M. *Lotus japonicus* E3 ligase SEVEN IN ABSENTIA4 destabilizes the symbiosis receptor-like kinase SYMRK and negatively regulates rhizobial infection. *Plant Cell*. 2012;24:1691–707.
25. Shimomura K, Nomura M, Tajima S, Kouchi H. LjnsRING, a novel RING finger protein, is required for symbiotic interactions between *Mesorhizobium loti* and *Lotus japonicus*. *Plant Cell Physiol*. 2006;47:1572–81.
26. Lu D, Lin W, Gao X, Wu S, Cheng C, Avila J, Heese A, Devarenne TP, He P, Shan L. Direct ubiquitination of pattern recognition receptor FLS2 attenuates plant innate immunity. *Science*. 2011;332:1439–42.
27. Liao D, Cao Y, Sun X, Espinoza C, Nguyen C, Liang Y, Stacey G. *Arabidopsis* E3 ubiquitin ligase PLANT U-BOX13 (PUB13) regulates chitin receptor LYSIN MOTIF RECEPTOR KINASE5 (LYK5) protein abundance. *New Phytol*. 2017;214:1646–56.
28. Li W, Ahn IP, Ning Y, Park CH, Zeng L, Whitehill JGA, Lu H, Zhao Q, Ding B, Xie Q, et al. The U-box/ARM E3 ligase PUB13 regulates cell death, defense, and flowering time in *Arabidopsis*. *Plant Physiol*. 2012;159:239–50.
29. Fukai E, Soyano T, Umehara Y, Nakayama S, Hirakawa H, Tabata S, Sato S, Hayashi M. Establishment of a *Lotus japonicus* gene tagging population using the exon-targeting endogenous retrotransposon LORE1. *Plant J*. 2012;69:720–30.
30. Urbański DF, Malolepszy A, Stougaard J, Andersen SU. Genome-wide LORE1 retrotransposon mutagenesis and high-throughput insertion detection in *Lotus japonicus*. *Plant J*. 2012;69:731–41.
31. Krusell L, Madsen LH, Sato S, Aubert G, Genua A, Szczygłowski K, Duc G, Kaneko T, Tabata S, de Bruijn F, Pajuelo E, Sandal N, Stougaard J. Shoot control of root development and nodulation is mediated by a receptor-like kinase. *Nature*. 2002;420:422–6.
32. Nishimura R, Hayashi M, Wu GJ, Kouchi H, Imaizumi-Anraku H, Murakami Y, Kawasaki S, Akao S, Ohmori M, Nagasawa M, Harada K, Kawaguchi M. HAR1 mediates systemic regulation of symbiotic organ development. *Nature*. 2002;420:426–9.
33. Zhou J, Lu D, Xu G, Finlayson SA, He P, Shan L. The dominant negative ARM domain uncovers multiple functions of PUB13 in *Arabidopsis* immunity, flowering, and senescence. *J Exp Bot*. 2015;66:3353–66.
34. Chinchilla D, Shan L, He P, de Vries S, Kemmerling B. One for all, the receptor-associated kinase BAK1. *Trends Plant Sci*. 2009;14:535–41.
35. Krusell L, Sato N, Fukuhara I, Koch BE, Grossmann C, Okamoto S, Okakura E, Otsubo Y, Aubert G, Nakagawa T, Sato S, Tabata S, Duc G, Parniske M, Wang TL, Kawaguchi M, Stougaard J. The *Clavata2* genes of pea and *Lotus japonicus* affect autoregulation of nodulation. *Plant J*. 2011;65:861–71.
36. Rasmussen SR, Führtbauer W, Novero M, Volpe V, Malkov N, Genre A, Bonfante P, Stougaard J, Radutoiu S. Intracellular colonization by arbuscular mycorrhizal fungi triggers induction of a lipochitoooligosaccharide receptor. *Sci Rep*. 2016;6:29733.
37. Malolepszy A, Urbański DF, James EK, Sandal N, Isono E, Stougaard J, Andersen SU. The deubiquitinating enzyme AMSH1 is required for rhizobial infection and nodule organogenesis in *Lotus japonicus*. *Plant J*. 2015;83:719–31.
38. Antignani V, Klocko AL, Bak G, Chandrasekaran SD, Dunivin T, Nielsen E. Recruitment of PLANT U-BOX13 and the P4Kb1/b2 phosphatidylinositol-4 kinases by the small GTPase RabA4B plays important roles during salicylic acid-mediated plant defense signalling in *Arabidopsis*. *Plant Cell*. 2015;27:243–61.
39. Yang CW, González-Lamothe R, Ewan RA, Rowland O, Yoshioka H, Shenton M, Ye H, O'Donnell E, Jones JD, Sadanandom A. The E3 ubiquitin ligase activity of *Arabidopsis* PLANT U-BOX17 and its functional tobacco homolog ACRE276 are required for cell death and defense. *Plant Cell*. 2006;18:1084–98.
40. Wang C, Xu X, Hong Z, Feng Y, Zhang Z. Involvement of ROP6 and clathrin in nodulation factor signalling. *Plant Signal Behav*. 2015b;10:e1033127.
41. Wang C, Zhu M, Duan L, Yu H, Chang X, Li L, Kang H, Feng Y, Zhu H, Hong Z, Zhang Z. *Lotus japonicus* clathrin heavy chain 1 is associated with rho-like GTPase ROP6 and involved in nodule formation. *Plant Physiol*. 2015a;167:1497–510.
42. Ke D, Fang Q, Chen C, Zhu H, Chen T, Chang X, Yuan S, Kang H, Ma L, Hong Z, Zhang Z. The small GTPase ROP6 interacts with NFR5 and is involved in nodule formation in *Lotus japonicus*. *Plant Physiol*. 2012;159:131–43.
43. Handberg K, Stougaard J. *Lotus japonicus*, an autogamous, diploid legume species for classical and molecular genetics. *Plant J*. 1992;2:487–96.
44. Broughton WJ, Dilworth MJ. Control of leghaemoglobin synthesis in snake beans. *Biochem J*. 1971;125:1075–80.
45. Kawaharada Y, Kelly S, Nielsen MW, Hjuler CT, Gysel K, Muszyński A, Carlson RW, Thygesen MB, Sandal N, Asmussen MH, et al. Receptor-mediated exopolysaccharide perception controls bacterial infection. *Nature*. 2015;523:308–12.
46. Tanou G, Minas IS, Karagiannis E, Tsikou D, Audebert S, Papadopoulou KK, Molassiotis A. The impact of sodium nitroprusside and ozone in kiwifruit ripening physiology, a combined gene and protein expression profiling approach. *Ann Bot*. 2015;116:649–62.

47. Weber E, Engler C, Gruetzner R, Werner S, Marillonnet S. A modular cloning system for standardized assembly of multigene constructs. *PLoS One*. 2011; 6:e16765.
48. Stougaard J, Abildsten D, Marcker KA. The *Agrobacterium rhizogenes* pRi TL-DNA segment as a gene vector system for transformation of plants. *Mol Gen Genomics*. 1987;207:251–5.
49. Hansen J, Jørgensen JE, Stougaard J, Marcker K. Hairy roots - a short cut to transgenic root nodules. *Plant Cell Rep*. 1989;8:12–5.
50. Vitha S, Benes K, Phillips JP, Gartland KM. Histochemical GUS analysis. *Methods Mol Biol*. 1995;44:185–93.
51. Madsen EB, Antolín-Llovera M, Grossmann C, Ye J, Vieweg S, Broghammer A, Krusell L, Radutoiu S, Jensen ON, Stougaard J, Parniske M. Autophosphorylation is essential for the in vivo function of the *Lotus japonicus* nod factor receptor 1 and receptor-mediated signalling in cooperation with nod factor receptor 5. *Plant J*. 2011;65:404–17.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

